

Insertion/Deletion Polymorphism of Angiotensin Converting Enzyme Gene in Kawasaki Disease

Polymorphism of angiotensin converting enzyme (ACE) gene is reported to be associated with ischemic heart disease, hypertrophic cardiomyopathy, and idiopathic dilated cardiomyopathy. In this study, we investigated the relationship between Kawasaki disease and insertion/deletion polymorphism of ACE gene. Fifty five Kawasaki disease patients and 43 healthy children were enrolled. ACE genotype was evaluated from each of the subjects' DNA fragments through polymerase chain reaction (PCR). Frequencies of ACE genotypes (DD, ID, II) were 12.7%, 60.0%, 27.3% in Kawasaki group, and 41.9%, 30.2%, 27.9% in control group respectively, indicating low rate of DD and high rate of ID genotype among Kawasaki patients ($p<0.01$). Comparing allelic (I, D) frequencies, I allele was more prevalent in Kawasaki group than in control group (57.3% vs. 43.0%, $p<0.05$). In Kawasaki group, both genotype and allelic frequencies were not statistically different between those with coronary dilatations and those without. ACE gene I/D polymorphism is thought to be associated with Kawasaki disease but not with the development of coronary dilatations.

Key Words : Mucocutaneous Lymph Node Syndrome; Kawasaki Disease; Peptidyl-Dipeptidase A; Angiotensin Converting Enzyme; Polymorphism, Genetic

Yoon Hee Shim, Hae Soon Kim,
Sejung Sohn, Young Mi Hong

Department of Pediatrics, College of Medicine, Ewha Womans University, Seoul, Korea

Received : 8 June 2005

Accepted : 29 September 2005

Address for correspondence

Hae Soon Kim, M.D.
Department of Pediatrics, Ewha Womans University
Mokdong Hospital, 911-1 Mok-dong, Yangcheon-gu,
Seoul 158-710, Korea
Tel : +82-2-2650-5569, Fax : +82-2-2653-3718
E-mail : hyesk@ewha.ac.kr

INTRODUCTION

Kawasaki disease is an acute systemic vasculitis and its diagnosis is made on clinical features. Main complication of the disease is coronary artery lesion that may result in myocardial infarction or sudden death. Coronary artery aneurysms or ectasia develop in approximately 15-25% of untreated children (1). Since the intravenous immune globulin (IVIG) and aspirin therapy have been introduced, its mortality rate has decreased to 0.1%, but cardiac sequelae continues to occur in about 13% of Kawasaki disease patients (2).

The etiology of Kawasaki disease is largely unknown despite the various suggested hypotheses. Based on epidemiologic and clinical manifestations, it is thought that Kawasaki disease is caused by some infectious agents (3-10). Hypercytokinemia and hyperchemokinemia have also been observed and are thought to cause vascular injuries by inflammatory reaction and immunologic activation (11). Genetic factors are also thought to have influences on the development and progress of Kawasaki disease (12, 13). Up to present days, Kawasaki disease is thought to be an infectious disease manifested by immunologic reaction in genetically susceptible person.

Angiotensin converting enzyme (ACE) breaks down the potent vasodilator, bradykinin to its inactive metabolite and catalyzes angiotensin I to angiotensin II. Angiotensin II promotes hyperplasia and hypertrophy of vascular smooth muscle cells, induces the production of proinflammatory cytokines

and causes endothelial dysfunction by free radical generation. By playing an important role in cardiovascular regulatory system, ACE gene has been proposed to be associated with various cardiovascular diseases, such as ischemic heart disease, hypertrophic cardiomyopathy, idiopathic dilated cardiomyopathy, and vascular hypertrophy (14). Among the different ACE genetic loci, the insertion/deletion polymorphism coded within intron 16 has been studied in many literature to find the association with such diseases.

However, there has been few studies regarding the association between the polymorphism of ACE gene and Kawasaki disease. The present study investigates whether the I/D polymorphism of ACE gene (DD, ID, II) is associated with the prevalence and severity of Kawasaki disease among Korean pediatric populations.

MATERIALS AND METHODS

Study subjects

Fifty five Kawasaki patients (mean age 28.2 ± 25.2 months) diagnosed at Ewha Womans University Mokdong Hospital from January 2001 to June 2003, and 43 healthy children (mean age 28.5 ± 17.2 months) were enrolled. Kawasaki disease was diagnosed by its clinical features, that is fever lasting for at least 5 days, accompanied by 4 of the 5 classical

signs: 1) bilateral bulbar conjunctival injection; 2) pharynx, injected and/or dry fissured lips, strawberry tongue; 3) changes of the peripheral extremities in the acute phase or periungual desquamation in the subacute phase; 4) nonvesicular rash; 5) cervical adenopathy, ≥ 1.5 cm. The diagnosis was made if he/she had typical manifestations even before less than 5 febrile days of illness. All patients were treated with immunoglobulin (2 g/kg) on the day of diagnosis. High dose (50 mg/kg/day) aspirin was given from the day of diagnosis and its dosage was changed to 5 mg/kg/day after 2 nonfebrile consecutive days.

Two dimensional echocardiography was done to evaluate cardiac complications. Coronary arterial lesion was defined as following: 1) inner diameter that is >3 mm in children <5 yrs old and >4 mm in children ≥ 5 yrs old; 2) internal diameter of a segment ≥ 1.5 times that of an adjacent segment; or 3) lumen with irregular surface. Among these patients, 17 showed coronary dilatations and 38 did not.

Informed consent was obtained from his/her parents prior to the participation in the study. The study was approved by the hospital's ethics committee.

ACE genotyping

Each of the subjects' DNA was extracted from whole blood at the time of diagnosis using QIAamp DNA Blood Mini Kit (Gene Company LTD., Chai Wan, Hong Kong). DNA fragments were amplified through polymerase chain reaction (PCR), which was carried out in a total volume of 10 μ L containing 50 ng of genomic DNA, 200 mM dNTPs, 0.3 mM/mL of each primers (5'-CTGGAGACCACTCCCATCCTTCT); (5'-GATGTGGCCATCACATTCTCAGAT) in PCR buffer with 0.5 units *Taq* DNA polymerase (Takara, Shiga, Japan). After the initial denaturation step (10 min at 95°C), 35 cycles

Table 1. Demographic characteristics of Kawasaki disease patients and control group

	Kawasaki disease			Control
	Normal Co A.	Dilated Co A.	Total	
Number	38	17	55	43
Age (month)	26.8 \pm 19.6	31.8 \pm 36.3	28.2 \pm 25.2	28.5 \pm 17.2
Sex (M:F)	20:18	8:9	28:27	30:13

$p > 0.05$. Co A., coronary artery.

Table 3. Deletion/insertion allelic prevalence of angiotensin converting enzyme gene in Kawasaki disease patients and control group

No. (%)	Allele		Total
	D	I	
Control	49 (57.0)	37 (43.0)	86 (100.0)
Kawasaki	47 (42.7)	63 (57.3)	110 (100.0)

$*p < 0.05$. odds ratio=1.78.

were repeated for 30 sec at 94°C, 30 sec at 52°C, 90 sec at 72°C, and 5 min at 72°C. DNA fragments were then separated by electrophoresis on 2.5% agarose gel.

Statistical analysis

Student's t-test was used to compare the demographic characteristics between the two groups. Chi square test and Fisher exact test were performed to compare the genotype and allelic frequencies between the two groups. Its frequencies were compared by odds ratio. p value less than 0.05 was considered significant.

RESULTS

The mean age of Kawasaki group (28 boys and 27 girls) was 28.2 ± 25.2 months, and that of the control group (30 boys and 13 girls) was 28.5 ± 17.2 months. Among 55 Kawasaki disease patients, coronary dilatation was observed in 17 patients (8 boys and 9 girls, Table 1)

Frequencies of ACE genotypes (DD, ID, II) were 12.7%, 60.0%, 27.3% in Kawasaki group, and 41.9%, 30.2%, 27.9% in control group respectively, indicating low rate of DD genotype ($p < 0.01$, odds ratio=0.2) and high rate of ID genotype ($p < 0.01$, odds ratio=3.3) among Kawasaki patients (Table 2).

Comparing allelic (I, D) frequencies, I allele was more prevalent in Kawasaki group than in control group (57.3% vs. 43.0%, $p < 0.05$, odds ratio=1.78, Table 3).

In Kawasaki group, both genotype (DD, ID, II) and allelic frequencies were not statistically different between those with coronary dilatations and those without (Table 4, 5).

Table 2. Prevalence of angiotensin converting enzyme genotype in Kawasaki disease patients and control group

No. (%)	ACE genotype			Total
	DD*	ID†	II	
Control	18 (41.9)	13 (30.2)	12 (27.9)	43 (100.0)
Kawasaki	7 (12.7)	33 (60.0)	15 (27.3)	55 (100.0)

* $p < 0.01$; odds ratio=0.2; † $p < 0.01$; odds ratio=3.3. ACE, angiotensin converting enzyme.

Table 4. Angiotensin converting enzyme genotype of Kawasaki patients with and without coronary dilatation

No. (%)	ACE genotype			Total
	DD	ID	II	
Normal Co A.	5 (13.1)	24 (63.2)	9 (23.7)	38 (100.0)
Dilated Co A.	2 (11.8)	9 (52.9)	6 (35.3)	17 (100.0)
Total	7 (12.7)	33 (60.0)	15 (27.3)	55 (100.0)

* $p > 0.05$. Co A., coronary artery; ACE, angiotensin converting enzyme.

Table 5. Deletion/insertion allelic prevalence of angiotensin converting enzyme gene of Kawasaki patients with and without coronary dilatation

No. (%)	Allele		Total
	D	I	
Normal Co A.	34 (44.7)	42 (55.3)	76 (100.0)
Dilated Co A.	13 (41.2)	21 (58.8)	34 (100.0)
Total	47 (43.6)	63 (56.4)	110 (100.0)

* $p > 0.05$. Co A., coronary artery.

DISCUSSION

The *ACE* gene is localized on chromosome 17q23 and is characterized by a major insertion/deletion polymorphism consisting of the presence or absence of a 287-base pair *Alu* repeat sequence within intron 16 (15). Angiotensin converting enzyme is an ectoenzyme found on the external surface of the endothelial and epithelial cell membranes. It enhances the synthesis of angiotensin-II, that promotes proliferation, migration, and hypertrophy of vascular smooth muscle cells. Angiotensin II also induces the production of proinflammatory cytokines and matrix metalloproteinases (16, 17). Moreover, the increased free radical generation by angiotensin-II contributes to endothelial dysfunction (18, 19).

In humans, the *ACE* activity is partly under genetic control. It is suggested that about half of the interindividual difference in *ACE* levels may be accounted for its polymorphism (19, 20). It is reported that mean *ACE* levels were lowest for II homozygotes, highest for DD homozygotes, and intermediate for ID heterozygotes (18, 21). Danser et al. explained the higher *ACE* levels observed in subjects with D allele than those with the II genotype by the sequence harbored in the insert of *ACE* gene. This sequence was said to be very similar to a silencer element (22). But it is not clear whether increased levels of *ACE* actually affects the levels of angiotensin II because the renin-angiotensin system is regulated by feedback mechanism (22).

Angiotensin converting enzyme is mainly produced by vascular endothelial cells (23). In Kawasaki disease, the associated endothelial cell damage subsequently lowers the *ACE* level. It is reported in some literature (24, 25) that serum *ACE* levels are significantly attenuated during the acute phase, and recovered during the convalescent phase of Kawasaki disease.

Slowik et al. (26) reported that the II genotype of *ACE* gene contributes to vascular dilatation at the site of aneurysm by 1) increased bradykinin activity, 2) another polymorphism responsible for vascular dilatation that is in linkage disequilibrium with *ACE* I/D polymorphism, 3) degeneration of endothelial cells, or 4) lack of vascular remodeling.

There is pathological difference between adult coronary artery disease (CAD) and that caused by Kawasaki disease. CAD caused by Kawasaki disease is characterized by vascular intimal thickening, whereas the adult CAD is characteri-

zed by atherosclerotic lesions initiated by atheroma and plaque formation. Thus, the pathophysiologic mechanism of myocardial ischemic development is also different; in adult CAD, plaque rupture and thrombus formation plays the important role, whereas in Kawasaki disease, coronary arterial narrowing by intimal hyperplasia is responsible (24).

In this study, the ID genotype was more prevalent (59.3 vs. 30.2%, $p < 0.01$), and the DD genotype (12.9 vs. 41.9%, $p < 0.01$) was less prevalent in Kawasaki group than in control group. And I allele was more prevalent in Kawasaki group than in control group. However, both genotype (DD, ID, II) and allelic frequencies were not statistically different between Kawasaki disease patients with coronary dilatations and those without. Therefore, *ACE* gene I/D polymorphism was thought to be associated with the prevalence of Kawasaki disease but not with the development of coronary lesions. These results are similar to the report by Wu et al. (13) that the DD genotype was present in lower frequency among Kawasaki patients and *ACE* polymorphism was not associated with coronary aneurysmal formation. These results are different from the study by Takeuchi et al. (27) that the II genotype of *ACE* gene is more prevalent in Kawasaki disease and those with coronary aneurysm. These inconsistent results may be accounted for different ethnic traits of the individual population.

In conclusion, the ID genotype is present in a significantly higher frequency and DD in lower frequency among Kawasaki disease patients than in control subjects. In addition, there are no significant association between *ACE* I/D polymorphism and the coronary artery aneurysm formation in Kawasaki disease patients. Because the study groups are relatively small size in number, it is difficult to generalize these results. In this study, the coronary arterial dilatation was observed in 17 out of 55 Kawasaki patients. This is higher than that in previously reported literature. It can be accounted for either 1) longer duration from the onset of illness to the infusion of immunoglobulin, 2) other factors that may affect the development of coronary arterial lesions in Kawasaki disease, such as different ethnicity or 3) selection bias caused by limited number of enrolled institution and small number of patients. IVIG is recommended to be given within the first 10 days of illness and, if possible, within 7 days of illness. Since the duration of febrile days before the diagnosis and the IVIG infusion were 4.6 ± 1.6 days (range, 2–8 days) in our study groups, the first explanation seems less likely.

Further study is required to clarify the association.

REFERENCES

1. Newburger JW, Fulton DR. Kawasaki disease. *Curr Opin Pediatr* 2004; 16: 508-14.
2. Yanagawa H, Yashiro M, Nakamura Y. 14th nationwide survey on Kawasaki disease. *J Pediatr Practice* 1998; 61: 406-20.
3. Shulman ST, Rowley AH. Does Kawasaki disease have a retroviral

- aetiology? *Lancet* 1986; 2: 545-6.
4. Esper F, Shapiro ED, Weibel C, Ferguson D, Landry ML, Kahn JS. Association between a novel human Coronavirus and Kawasaki disease. *J Infect Dis* 2005; 191: 499-502.
 5. Chou CT, Chang JS, Ooi SE, Huo AP, Chang SJ, Chang HN, Tsai CY. Serum anti-Yersinia antibody in Chinese patients with Kawasaki disease. *Arch Med Res* 2005; 36: 14-8.
 6. Hagiwara K, Komura H, Kishi P, Kaji T, Yoshida T. Isolation of human herpesvirus-6 from an infant with Kawasaki disease. *Eur J Pediatr* 1992; 151: 867-8.
 7. Baba K, Takeda N, Tanaka M. Cases of Yersinia pseudotuberculosis infection having diagnostic criteria of Kawasaki disease. *Contrib Microbiol Immunol* 1991; 12: 292-6.
 8. Lee DH, Huang HP. Kawasaki disease associated with chickenpox: report of two sibling cases. *Acta Paediatr Taiwan* 2004; 45: 94-6.
 9. Kang JH, Lee JS. The role of EBV in the development of Kawasaki disease. 5th West-Pacific allergy symposium 1997; 241-4.
 10. Rocholl C, Gerber K, Daly J, Pavia AT, Byington CL. Adenoviral infections in children: the impact of rapid diagnosis. *Pediatrics* 2004; 113: e51-6.
 11. Kim DS. Immunologic aspect in Kawasaki disease. *J Korean Pediatr Cardiol Soc* 2003; 7: 306-10.
 12. Rowley AH, Shulman ST. Kawasaki syndrome. *Pediatric Clin N Am* 1999; 46: 313-29.
 13. Wu SF, Chang JS, Peng CT, Shi YR, Tsai FJ. Polymorphism of angiotensin-I converting enzyme gene and Kawasaki disease. *Pediatr Cardiol* 2004; 25: 529-33.
 14. Hubert C, Houot AM, Corvol P, Soubrier F. Structure of angiotensin I-converting enzyme gene. Two alternate promoters correspond to evolutionary steps of a duplicated gene. *J Biol Chem* 1991; 266: 15377-83.
 15. Manning MW, Cassis LA, Daugherty A. Differential effects of doxycycline, a broad-spectrum matrix metalloproteinase inhibitor, on angiotensin II-induced atherosclerosis and abdominal aortic aneurysms. *Arterioscler Thromb Vasc Biol* 2003; 23: 483-8.
 16. Lamblin N, Hermant X, Lablanche JM, Helbecque N, Amouyel P, Bauters C. Angiotensin converting enzyme and angiotensin II type I receptor polymorphisms in patients with coronary aneurysms. *Thromb J* 2003; 1: 5-11.
 17. Prasad A, Narayanan S, Waclawiw MA, Epstein N, Quyyumi AA. The insertion/deletion polymorphism of the angiotensin-converting enzyme gene determines coronary vascular tone and nitric oxide activity. *J Am Coll Cardiol* 2000; 36: 1579-86.
 18. Scheer WD, Boudreau DA, Hixson JE, McGill HC, Newman WP 3rd, Tracy RE, Zieske AW, Stong JP. ACE insert/delete polymorphism and atherosclerosis. *Atherosclerosis* 2005; 178: 241-7.
 19. Rigat B, Hubert C, Alhenc-Gelas F, Cambien F, Corvol P, Soubrier F. An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *J Clin Invest* 1990; 86: 1343-6.
 20. Tiret L, Rigat B, Visvikis S, Breda C, Corvol P, Cambien F, Soubrier F. Evidence, from combined segregation and linkage analysis, that a variant of the angiotensin I converting enzyme (ACE) gene controls plasma ACE levels. *Am J Hum Genet* 1992; 51: 197-205.
 21. Frossard PM, Lestringant GG, Obineche EN, Hill SH. The angiotensin-converting enzyme (ACE) gene insertion/deletion dimorphism tracks with higher serum ACE activities in both younger and older subjects. *Ann Saudi Med* 1998; 18: 289-392.
 22. Danser AH, Schunkert H. Renin-angiotensin system gene polymorphism: potential mechanisms for their association with cardiovascular diseases. *Eur J Pharmacol* 2000; 410: 303-16.
 23. Caldwell PR, Segal BC, Hsu KC, Das M, Softer RL. Angiotensin converting enzyme: vascular endothelial localization. *Science* 1976; 191: 1050-1.
 24. Falcini F, Generini S, Pigone A, Leoncini G, Cimaz R, Partsch G, Matucci-Cerinic M. Are angiotensin converting enzyme and von Willebrand factor circulating levels useful surrogate parameters to monitor disease activity in Kawasaki disease. *Endothelium* 1999; 6: 209-15.
 25. Fukazawa R, Sonobe T, Hamamoto K, Hamaoka K, Sakata K, Asano T, Imai T, Kamisago M, Ohkubo T, Uchikoba Y, Ikegami E, Watanabe M, Ogawa S. Possible synergic effect of angiotensin-I converting enzyme gene insertion/deletion polymorphism and angiotensin-II type-1 receptor 1166A/C gene polymorphism on ischemic heart disease in patients with Kawasaki disease. *Pediatr Res* 2004; 56: 597-601.
 26. Slowik A, Borratynsky A, Pera J, Betlej M, Dziedzic T, Krzyszkowski T, Czepko R, Figlewicz DA, Szczudlik A. II genotype of the angiotensin-converting enzyme gene increases the risk for subarachnoid hemorrhage from ruptured aneurysm. *Stroke* 2004; 35: 1594-97.
 27. Takeuchi K, Yamamoto K, Kataoka S, Kakihara T, Tanaki A, Sato S, Uchiyama M. High incidence of angiotensin I converting enzyme genotype II in Kawasaki disease patients with coronary aneurysm. *Eur J Pediatr* 1997; 156: 266-8.