



What is the minimal supersymmetric Standard Model from F-theory?

Kang-Sin Choi ^{a,b,*}

^a Scranton College, Ewha Womans University, Seoul 120-750, Republic of Korea

^b Korea Institute for Advanced Study, Seoul 130-722, Republic of Korea

ARTICLE INFO

Article history:

Received 30 May 2012

Received in revised form 12 September 2012

Accepted 9 October 2012

Available online 11 October 2012

Editor: M. Cvetič

ABSTRACT

We construct gauge theory of $SU(3) \times SU(2) \times U(1)$ by spectral cover from F-theory and ask how the Standard Model is extended under minimal assumptions on Higgs sector. For the requirement on different numbers between Higgs pairs and matter generations (respectively one and three) distinguished by R-parity, we choose a universal G-flux obeying $SO(10)$ but slightly breaking E_6 unification relation. This condition forces distinction between up and down Higgs fields, suppression of proton decay operators up to dimension five, and existence and dynamics of a singlet related to μ -parameter.

© 2012 Elsevier B.V. Open access under CC BY license.

1. Introduction

We explore a supersymmetric extension of the Standard Model (SM) from F-theory, under certain minimal assumptions on Higgs sector. Construction from F-theory, admitting dual $E_8 \times E_8$ heterotic string, naturally yields a realistic Grand Unified Theory (GUT) of gauge group along E_n series, including SM itself [1,2]. In string derived models, however, such unification relation is so strong that it has been very difficult to understand the nature of Higgs doublet in this context, namely how to embed it to a larger GUT representation and why its observed number should be different from that of quark and lepton generations. The main result of this Letter is that F-theory can control such features, implying some nontrivial phenomenological consequences. For example it gives us understanding on how can we distinguish up and down-type Higgs fields and what are the properties of the μ -parameter in the Minimal Supersymmetric Standard Model (MSSM).

We first build $SU(3) \times SU(2) \times U(1)_Y$ gauge group, without aid of an intermediate Grand Unification. By specifying a spectral cover of structure group $S[U(5) \times U(1)_Y]$, its commutant in E_8 survives as the SM gauge group [4,5]. The spectral cover is a systematic way to construct (poly)stable vector bundle in dual heterotic string, if the compact manifold admits elliptic fibration with a (usually called zero) section [6]. Although the desired spectral cover is obtained by tuning parameters of an $SU(6)$ cover [7–9], the existence of the $U(1)_Y$ gauge group is not guaranteed until the following two requirements are met. First, elliptic fiber of heterotic string admits

more global section(s) than the zero section, since monodromy should not mix the single cover for $U(1)_Y$ from extension to non-abelian structure group [10,11]. This we do by tuning elliptic fiber as well [12]. Second, the corresponding gauge boson should not acquire mass by Stückelberg mechanism, which we evade by not turning on G-flux along this direction.

To obtain chiral spectrum in four dimension, we also have to turn on so-called G-flux [13,14]. It is important to note the unique feature of F-theory that the unbroken group is solely determined by the spectral cover and G-flux only affects the number of zero modes. Thus, as long as the spectral cover has the structure group $S[U(5) \times U(1)_Y]$ the unbroken group is the SM group. To distinguish Higgs from lepton doublet in supersymmetric model by R-parity and also to impose different number of Higgs from that of the unified matter multiplets, it will turn out that the structure group of spectral cover is singled out to be $S[U(3)_\perp \times U(1) \times U(1) \times U(1)]$.

If the structure group is semi-simple and possibly plus abelian, we can partly turn on the G-flux on a subgroup. For example, turning on G-flux on $SU(3)_\perp$ group, the resulting number of generation obeys the unification relation of the commutant group E_6 , predicting the same number of fields belonging to **27** multiplet of E_6 , thus the number of Higgs doublet should be the same as that of quark generations. This relation can be relaxed, on the other hand, if G-flux is on $SU(4)$, giving unification relation of $SO(10)$. This is attempted in the previous work [4], but the number of Higgs doublet is also totally determined to be undesirable one. To control them differently we turn on two different G-fluxes along its subgroup $S[U(3)_\perp \times U(1)] \subset SU(4)$ with one more free parameter. Since it does not obey E_6 unification the number of Higgs pair can be different to that of matter quarks, to be three and one, respectively, adjusted by $U(1)$ flux strength. The entire flux still does not

* Correspondence to: Scranton College, Ewha Womans University, Seoul 120-750, Republic of Korea.

E-mail address: kangsin@ewha.ac.kr.

touch $SO(10)$ direction, the model possesses $SO(10)$ unification relation thus we have the same number of quarks and leptons, as well as that of right-hand neutrinos.

Finally, the four-dimensional interactions follow from gauge invariant terms of the higher dimensional effective Lagrangian by dimensional reduction [15,16]. The invariance under the various $U(1)$ groups from the above spectral cover plays the role of selection rule. The structure of these symmetries predicts aforementioned phenomenological features. Also, we analyze the vacuum configuration giving proper interactions evading nucleon decays.

2. Gauge group

The model is obtained from F-theory compactification on elliptic Calabi–Yau fourfold with a section, admitting heterotic dual. The dual heterotic string is compactified on elliptic Calabi–Yau threefold $Z \rightarrow B_2$ with a section, which is usually called as the zero section. To have a globally well-defined $U(1)$ used by the SM gauge group and its constructing spectral cover, we need *another global section* than the zero section on the fiber to parameterize the dual point to the line bundle of the $U(1)$ structure group [12]. Globally, this point will not be mixed by monodromy with other points parameterizing other spectral covers, as we move around the entire base B_2 . Let the canonical bundle of the base B_2 be K_{B_2} . We choose the coordinate of such point as (x_1, y_1) , which are global holomorphic sections $x \in \Gamma(B_2, O(K_{B_2}^{-2}))$ and $y \in \Gamma(B_2, O(K_{B_2}^{-3}))$, and the elliptic equation has a form

$$(y - y_1)(y + y_1) = (x - x_1)(x^2 + x_1x + x_1^2 + f) \quad (1)$$

where $f \in \Gamma(B_2, O(K_{B_2}^{-4}))$.

We construct the spectral cover for the structure group $S[U(5) \times U(1)_Y]$ as follows [4,5,12],

$$a_0 + a_2x + a_3y + a_4x^2 + a_5xy + a_6x^3 = 0, \quad (2)$$

with tuning of parameters

$$\begin{aligned} a_0 &= d_0, & a_2 &= d_2 + b_1d_1, & a_3 &= d_3 + b_1d_2, \\ a_4 &= d_4 + b_1d_3, & a_5 &= d_5 + b_1d_4, & a_6 &= b_1d_5, \end{aligned} \quad (3)$$

with the constraint $d_1 + b_1d_0 = 0$. Here $a_m \in \Gamma(B_2, O(K_{B_2}^m))$ are globally defined and no approximation, e.g. of Higgs bundle type is used. In addition, to guarantee the existence of a global section with holomorphic parameters, we need further factorization condition [12]

$$f = b_1^2F, \quad g = -b_0^2F, \quad d_0 = b_0d, \quad d_1 = -b_1d, \quad (4)$$

where the topological properties of d and F can be deduced from those of f , g , b_0 , b_1 . Since the global section $(x_1, y_1) = (b_0^2/b_1^2, \pm b_0^3/b_1^3)$ is on this spectral cover (2), the coordinate values will be expressed in terms of the parameters b_1 and d_m . We can take an analogy of Higgs bundle for large x and y to plug well-known solution so far (but we do not stick to it since our description is valid for all x and y as long as the stable degeneration limit is valid): Each coefficient d_m , parameterizing the positions of the covers, is related to the elementary symmetric polynomial of degree m , out of weights of the fundamental representations $\mathbf{5}_1 + \mathbf{1}_{-5}$ of the $S[U(5) \times U(1)_Y]$. The surviving group on B_2 is the commutant, the SM group $SU(3) \times SU(2) \times U(1)_Y$. This a sufficient specification, so that it provides the information on the unbroken gauge group [19].

In the stable degeneration limit [6,14], we can convert Eqs. (1) and (2) into the singularity equation corresponding to the SM group

Table 1

Matter contents identified by $SU(3) \times SU(2) \times SU(3)_\perp \times U(1)_Y \times U(1)_X \times U(1)_Z$ quantum numbers. All the indices take different value in $S_3 = \{1, 2, 3\}$. Later, the fields below middle line are decoupled and the charge conjugates of h_u^c and D_1^c will survive as zero modes.

Matter	Matter curve	Homology on B_2
$q(\mathbf{3}, \mathbf{2}; \mathbf{3})_{\frac{1}{6}, 1, 1}$	$\Pi t_i \rightarrow 0$	$\eta - 3c_1$
$u^c(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{3})_{-\frac{2}{3}, 1, 1}$	$\Pi(t_i + t_6) \rightarrow 0$	$\eta - 3c_1$
$e^c(\mathbf{1}, \mathbf{1}; \mathbf{3})_{1, 1, 1}$	$\Pi(t_i - t_6) \rightarrow 0$	$\eta - 3c_1$
$d^c(\mathbf{3}, \mathbf{1}; \mathbf{3})_{\frac{1}{2}, -3, 1}$	$\Pi(t_i + t_5) \rightarrow 0$	$\eta - 3c_1$
$l(\mathbf{2}, \mathbf{1}; \mathbf{3})_{-\frac{1}{2}, -3, 1}$	$\Pi(t_i + t_5 + t_6) \rightarrow 0$	$\eta - 3c_1$
$v^c(\mathbf{1}, \mathbf{1}; \mathbf{3})_{0, 5, 1}$	$\Pi(t_i - t_5) \rightarrow 0$	$\eta - 3c_1$
$h_u^c(\mathbf{2}, \mathbf{1}; \bar{\mathbf{3}})_{\frac{1}{2}, 2, 2}$	$\Pi(t_i + t_j + t_6) \rightarrow 0$	$\eta - 3c_1$
$h_d(\mathbf{2}, \mathbf{1}; \mathbf{3})_{\frac{1}{2}, 2, -2}$	$\Pi(t_i + t_4 + t_6) \rightarrow 0$	$\eta - 3c_1$
$D_1^c(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{3})_{\frac{1}{3}, 2, 2}$	$\Pi(t_i + t_j) \rightarrow 0$	$\eta - 3c_1$
$\bar{D}_2(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{3})_{\frac{1}{3}, 2, -2}$	$\Pi(t_i + t_4) \rightarrow 0$	$\eta - 3c_1$
$S(\mathbf{1}, \mathbf{1}; \mathbf{3})_{0, 0, 4}$	$\Pi(t_i - t_4) \rightarrow 0$	$\eta - 3c_1$
$X(\mathbf{3}, \mathbf{2}; \mathbf{1})_{-\frac{5}{6}, 0, 0}$	$t_6 \rightarrow 0$	$-c_1$
$Y(\mathbf{3}, \mathbf{2}; \mathbf{1})_{\frac{1}{6}, -4, 0}$	$t_5 \rightarrow 0$	$-c_1$
$T^c(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1})_{-\frac{2}{3}, -4, 0}$	$t_5 + t_6 \rightarrow 0$	$-c_1$
$\Sigma(\mathbf{1}, \mathbf{1}; \mathbf{1})_{1, -4, 0}$	$t_5 - t_6 \rightarrow 0$	$-c_1$
$Q(\mathbf{3}, \mathbf{2}; \mathbf{1})_{\frac{1}{6}, 1, -3}$	$t_4 \rightarrow 0$	$-c_1$
$U^c(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1})_{-\frac{2}{3}, 1, -3}$	$t_4 + t_6 \rightarrow 0$	$-c_1$
$E^c(\mathbf{1}, \mathbf{1}; \mathbf{1})_{1, 1, -3}$	$t_4 - t_6 \rightarrow 0$	$-c_1$
$D^c(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1})_{\frac{1}{3}, -3, -3}$	$t_4 + t_5 \rightarrow 0$	$-c_1$
$L(\mathbf{2}, \mathbf{1}; \mathbf{1})_{-\frac{1}{2}, -3, -3}$	$t_4 + t_5 + t_6 \rightarrow 0$	$-c_1$
$N^c(\mathbf{1}, \mathbf{1}; \mathbf{1})_{0, 5, -3}$	$t_4 - t_5 \rightarrow 0$	$-c_1$

$$\begin{aligned} y^2 &= x^3 + (d_5 + d_4b_1)xy + (d_3 + d_2b_1)(b_1d_5 + z)yz \\ &\quad + (d_4 + d_3b_1)x^2z + (d_2 - b_1^2d)(b_1d_5 + z)xz^2 \\ &\quad + d(b_1d_5 + z)^2z^3 + b_1^2Fxz^4 - Fz^6, \end{aligned} \quad (5)$$

where x , y are affine coordinates of \mathbb{P}^2 and z is the coordinate of blown-up \mathbb{P}^1 in the stable degeneration [14]. Roughly, z is a normal coordinate to $B_2 = \{z = 0\}$ inside the base of elliptic fibration B in the F-theory side. At the discriminant locus of (5), we have the SM gauge group [5]. Referring to Tate's table [19], already (5) is a special form of the $SU(3)$ singularity whose parameters are tuned up to $\mathcal{O}(z^5)$. A change of coordinate $a_1b_5 + z \rightarrow z$ shows the other $SU(2)$ part is also special up to $\mathcal{O}(z^5)$. The $U(1)_Y$ part is the relative position between two linearly equivalent components. Its global existence depends on the terms in the last line of (5) although they look sub-leading contribution in z , otherwise we cannot have a monodromy-invariant two cycle harboring two-form related to $U(1)_Y$ [12]. The Calabi–Yau conditions require that the b_m are sections of $\eta - mc_1$, where $\eta = 6c_1(B_2) + c_1(N_{B_2/B})$ and $c_1 = c_1(B_2)$ are combinations of tangent and normal bundle to B_2 . The leading order locus of the discriminant in z coincides with B_2 .

The spectral cover should be further decomposed with smaller structure group, due to phenomenological requirements. We need to distinguish Higgs doublets from lepton doublets, having the same SM quantum numbers. The standard way is to introduce the matter parity, or its continuous version $U(1)_X$ with the charge being the baryon minus the lepton numbers. This is the commutant to $SU(5)$ inside $SO(10)$ GUT group along E_n series, hence a subgroup of the structure group. So we may decompose the spectral cover with $U(1)_X$. Shortly we will see, for the observed number of Higgs fields in four dimension, we need one more parameter from an extra $U(1)_Z$, so that the structure group should be factorized as

$$S[U(3)_\perp \times U(1)_Z \times U(1)_X \times U(1)_Y]. \quad (6)$$

The resulting spectral cover, respectively $C_3 \cup C_Z \cup C_X \cup C_Y$, is realized by further tuning $d_0 = f_0$, $d_1 = f_1 + e_1f_0$, $d_2 = f_2 + e_1f_1$, $d_3 = f_3 + e_1f_2$, $d_4 = f_4 + e_1f_3$, $d_5 = e_1f_4$ with the constraint

$f_0(b_1 + d_1 + e_1) + f_1 = 0$. In Z , their classes are respectively $C_3 = 3\sigma + \pi^*\eta$ and we have linear equivalence relations $C_Z \sim C_X \sim C_Y = \sigma$.

3. Matter contents

Since we admit heterotic duality, all four-dimensional fields comes from branching and auction of the adjoint **248** of E_8 [18]. Accordingly it branches into multiplets of $SU(3) \times SU(2) \times SU(3)_\perp \times U(1)_Y \times U(1)_X \times U(1)_Z$. The matter spectrum is summarized in Table 1. We identify the fields by charge assignments

$$\begin{aligned} Y: & \left(\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, -\frac{5}{6} \right), \\ X: & (1, 1, 1, 1, -4, 0), \\ Z: & (1, 1, 1, -3, 0, 0), \end{aligned} \quad (7)$$

in the basis $\{t_1, t_2, t_3, t_4, t_5, t_6\}$, the weight vectors **5**₁ + **1**₋₅ of the structure group $S[U(5) \times U(1)_Y]$. They are localized along curves, the projections of $C_a \cap \tau C_b$ or $C_a \cap C_b$, $a, b \in \{3, Z, X, Y\}$ on B_2 , where τ is involution flipping the orientation of the cover.

The identities of extra singlets v^c and S are understood as follows. The minimal anomaly free single chiral representation containing all the observed fermions of SM is **16** of $SO(10)$. It also contains one extra SM singlet v^c . Invariance under $SO(10)$ forms Dirac mass term for v^c with the SM lepton doublet, thus this is to be interpreted as right-handed neutrino. With the aid of supersymmetry (SUSY), Higgs bosons belong to a hypermultiplet and can be treated on equal footing as matter. Thus matter and Higgs pair (as well as colored Higgs pair) are unified to a single representation **27** of E_6 . Again it predicts another kind of singlet S , and the gauge invariance relates this to μ -parameter of SUSY [17]. So the matter contents and couplings naturally show a singlet extension of minimal supersymmetric Standard Model.

The field strengths along the Cartan direction come from the dimensional reduction of four-form field strength G of the dual M-theory and this induces vector bundle on the spectral cover [6]. Although the minimal $SU(4)$ G -flux preserves unification relation of its commutant $SO(10)$ in E_8 , the number of Higgs pairs turns out to be completely fixed to be twice the matter multiplicity [4]. Here we have one more parameter ζ , the trace part of the $U(3)_\perp \subset SU(4)$ vector bundle [20], to relax the condition. So we turn on a universal flux

$$\begin{aligned} \Gamma_3 &= \lambda(3\sigma - \pi_3^*(\eta - 3c_1)) + \frac{1}{3}\pi_3^*\zeta, \\ \Gamma_Z &= -\pi_Z^*\zeta, \quad \Gamma_Y = \Gamma_X = 0, \end{aligned} \quad (8)$$

where σ is the class for B_2 inside Z and π_Z , π_Z are projections from $U(3)_\perp$ and $U(1)_Z$ covers to B_2 , respectively. In the F-theory side, we can turn off other fluxes along $U(1)_Y$ or $U(1)_X$ directions, as long as the quantization condition for λ below is satisfied. However there is no corresponding picture in the heterotic side, since Fourier–Mukai transformation with zero flux on some of the covers does not make sense.

The number n_R of chiral R zero modes minus anti-chiral \bar{R} ones of the Dirac operator in Z is a topological number and counted by index theorem. It is simply given by the intersection between matter curve class and Poincaré dual of the G -flux, projected on B_2 [14,21]

$$n_R = \mathcal{P}_R \cap \Gamma, \quad (9)$$

where \mathcal{P}_R is the matter curve of the representation R and \cap denotes the intersection inside Z . Because of identical geometry of

spectral cover as in Refs. [22,23], and we refer to it for the calculation of matter curves

$$\begin{aligned} n_q &= n_{u^c} = n_{e^c} = n_{d^c} = n_l = n_{v^c} \\ &= (3\sigma + \eta) \cap \sigma \cap \left(\lambda(3\sigma_\infty - \eta) + \frac{1}{3}\zeta \right) + (-c_1) \cap \sigma \cap (-\zeta) \\ &= \left(-\lambda\eta + \frac{1}{3}\zeta \right) \cdot (\eta - 3c_1) + c_1 \cdot \zeta, \end{aligned} \quad (10)$$

$$\begin{aligned} n_{D_1} &= n_{h_u} \\ &= -(2\sigma + \eta) \cap (\eta - 3c_1) \cap \left(\lambda(3\sigma_\infty - \eta) + \frac{1}{3}\zeta \right) \\ &= \left(-\lambda\eta - \frac{2}{3}\zeta \right) \cdot (\eta - 3c_1), \end{aligned} \quad (11)$$

$$\begin{aligned} n_{\bar{D}_2} &= n_{h_d} \\ &= (3\sigma + \eta) \cap \sigma \cap \left(\lambda(3\sigma_\infty - \eta) + \frac{1}{3}\zeta - \zeta \right) \\ &= \left(-\lambda\eta - \frac{2}{3}\zeta \right) \cdot (\eta - 3c_1), \end{aligned} \quad (12)$$

$$n_X = n_Y = n_{T^c} = n_\Sigma = 0, \quad (13)$$

$$n_Q = n_{U^c} = n_{E^c} = n_{D^c} = n_L = n_{N^c} = -c_1 \cdot \zeta, \quad (14)$$

$$\begin{aligned} n_S &= (3\sigma + \eta) \cap \sigma \cap \left(\lambda(3\sigma_\infty - \eta) + \frac{1}{3}\zeta + \zeta \right) \\ &= \left(-\lambda\eta + \frac{4}{3}\zeta \right) \cdot (\eta - 3c_1). \end{aligned} \quad (15)$$

Here we omitted pullback and the dot product is for the divisors of B_2 . We defined $\sigma_\infty = \sigma + \pi^*c_1$. All the matter fields appearing here are those inside **27** multiplet of E_6 . Their multiplicities manifest the $SO(10)$ unification relation, predicting the same number of right-handed neutrinos. They are preserved because the G -flux is along $SU(3)_\perp \times U(1)_Z$ structure group. It is a nontrivial check that h_u and h_d gives the same number in (10) and (11), so that there is no anomaly in four-dimension.

The numbers of matter generations and Higgs pairs can be individually controlled, depending on the topological data on B_2 . We require three generations of matter and one pair of Higgs doublets

$$\lambda\eta \cdot (\eta - 3c_1) = -\frac{7}{3}, \quad \eta \cdot \zeta = 2, \quad c_1 \cdot \zeta = 0. \quad (16)$$

They are subject to quantization conditions $3(\frac{1}{2} + \lambda) \in \mathbb{Z}$, $(\frac{1}{2} - \lambda)\eta + (3\lambda + \frac{1}{2})c_1 + \frac{1}{3}\zeta \in H_2(S, \mathbb{Z})$ where λ is a nonnegative rational number. We find a solution $\lambda = \frac{1}{6}$, for which only an integral or half-integral λ is possible in the absence of $U(1)_Z$ flux ζ . The base as del Pezzo two surface with $\eta = 2H$, $\zeta = H - 3E_1$ do the job, where H is hyperplane divisor and E_1 is one of the exceptional divisor. This relation restricts the number of the SM neutral field S be five. In addition, because c_1 in (2) transforms as a section of $-c_1$, we have two scalar fields O and O' transforming as adjoints under $S[U(3) \times U(2)] \cong SU(3) \times SU(2) \times U(1)$, belonging to $H^{2,0}(B_2) + H^{0,1}(B_2)$ [13]. They will play an interesting role in vacuum configuration around the string scale M_s . The other E_8 serves as hidden sector and is completely decoupled in smooth compactification and it can serve as supersymmetry breaking sector. In the F-theory side, we can turn off other fluxes along $U(1)_Y$ or $U(1)_X$ directions, as long as the quantization condition for λ is satisfied.

4. Higgs sector and nucleon decay

The requirement of *one pair* of Higgs doublets fixed the factorization of spectral cover (6). It has the following phenomenological implications.

Firstly, it also distinguishes between *up and down* type Higgses. This is due to the structure of the $SU(3)_\perp$ monodromy S_3 [24], the permutation of the elements $\{t_1, t_2, t_3\}$. It is the natural Weyl group, without a special monodromy further selected by hand. In terms of the S_3 representations, the fields having the same quantum number of lepton doublet under the SM group are

$$\begin{aligned} l: & \{t_1 + t_5 + t_6, t_2 + t_5 + t_6, t_3 + t_5 + t_6\}, \\ h_u^c: & \{t_1 + t_2 + t_6, t_2 + t_3 + t_6, t_3 + t_1 + t_6\}, \\ h_d: & \{t_1 + t_4 + t_6, t_2 + t_4 + t_6, t_3 + t_4 + t_6\}, \\ L: & \{t_1 + t_5 + t_6, t_2 + t_5 + t_6, t_3 + t_5 + t_6\}. \end{aligned} \quad (17)$$

Effectively, the Higgs doublet is distinguished from the lepton doublet by an opposite matter parity or the $U(1)_X$. It also forbids bare (super)renormalizable lepton and/or baryon number violating operators $lh_u, ll e^c, lq d^c, u^c d^c e^c$. Further factorization ruins this one Higgs pair structure but we obtain three pairs of Higgses, so our factorization seems the unique for the $U(n)$ type spectral cover with universal flux.

Well known is that the matter parity and $U(1)_X$ alone cannot forbid dimension five proton decay operators such as $qqql$ and $u^c u^c d^c e^c$. However, the above structure group *forbids* these operators. For instance, $qqql$ is not allowed because of nonvanishing sum of the weights $(t_i) + (t_j) + (t_k) + (t_l + t_5 + t_6)$ and $u^c u^c d^c e^c$ is not because of $(t_i + t_6) + (t_j + t_6) + (t_k + t_5) + (t_l - t_6)$, required by $SU(3)_\perp$ invariance, since one of S_3 index should appear twice [25]. At the field theory level, this is also simply understood by invariance under $U(1)_Z$ [26].

Another prediction is the presence of an SM singlet field S . Surveying the quantum number, it belongs to **27** representation of E_6 , therefore, its interaction is restricted and we can calculate the corresponding terms. Because Higgs doublet and triplets are not simply vectorlike and up and down Higgses live on different matter curves, bare masses are forbidden by $SU(3)_\perp$ invariance. Instead we have a singlet extension to MSSM [28,29]. We can check that the only renormalizable superpotential for the surviving fields are (see also below)

$$\begin{aligned} qh_d u^c + qh_u d^c + lh_d e^c + lh_u v^c + Sh_u h_d + SD_1 \bar{D}_2 \\ + qqD_1 + u^c e^c D_1 + q \bar{D}_2 + v^c d^c D_1 + u^c d^c \bar{D}_2 \end{aligned} \quad (18)$$

omitting the flavor dependent coefficients. We expect the terms involving D_1 and D_2^c are all decoupled, yielding the μ -like term $Sh_u h_d$. Bare quadratic or cubic terms in S are not allowed by invariance under the $SU(3)_\perp$ and other $U(1)$'s. Induced higher order terms include $M_s^{-2} SS(Q \bar{D}_2 L + D^c N D_1 + U^c E^c D_1) + M_s^{-4} SSS(Q U^c E^c L + Q D^c N^c L)$ but they are to be suppressed by a string scale M_s . A Majorana mass for the v^c does not appear up to dimension five. There is an interesting room for this from Euclidean D3-brane or M5-brane instanton in F-theory [30], which might as well generate similar potential for S .

Since the Higgs fields also obey $SO(10)$ unification relation, we have as many colored Higgs pairs D_1, \bar{D}_2 as doublets. This doublet-triplet splitting problem should be solved by an effect evading the unification structure, close but below M_s . It is a possibility to consider a *vectorlike* extra generation of matter fields, without changing the Dirac indices. Using aforementioned $U(3)$ adjoint chiral super field O , there can be terms $\langle O \rangle D_1 D_1^c + \langle O \rangle \bar{D}_2 \bar{D}_2^c + M_O \text{tr } O^2 + \text{tr } O^3$ giving Dirac masses separately to D_1 ,

D_1^c and \bar{D}_2, \bar{D}_2^c pairs. Conventional gauge coupling unification requires heavy triplets, so do a large VEV $\langle O \rangle$ and a large M_O [32]. We can allow also vectorlike pair for the doublet, but in principle a similar $U(2)$ adjoint can give different masses. This seems like a flavor problem in the UV regime and more is to be understood. On the other hand, we expect a coupling $(\langle S \rangle + \mu_D) D_1 \bar{D}_2$ is generated, with a possible SUSY breaking effect μ_D [31]. The most strongly constrained nucleon decay operator is $qqql$, whose coefficient has upper bound $10^{-5} M_p^{-1}$ [33]. At low energy scale, integrating out heavy fields, qqD_1 and $q \bar{D}_2$ may induce an operator $(\langle S \rangle + \mu_D) / M_D^{-2} qqql$ up to geometric suppression factor. Once forbidden at the tree-level, it is also known that the induced operators are highly suppressed, probably explained by worldsheet instanton contribution [27]. A possible mixing from bare mass term $d^c d$ does not change this eigenvalue. The same argument goes to other induced operators for nucleon decay.

5. Anomalous $U(1)$

We check the G -flux contribution to D -term for each $U(1)$ using type IIB string limit [34], where we have Ramond–Ramond four-form field C_4 in low energy. Its Kaluza–Klein expansion along a harmonic two-form $w_2 \in H^{1,1}(B_2, \mathbb{Z})$ has a form $C_4 = C_2 \wedge \omega_2$, yielding the interaction $\text{tr} t_Q^2 \int_{M^4} F_Q \wedge C_2 \int_{B_2} i^* \omega_2 \wedge (F_Q)$ from Chern–Simons interactions and here F_Q , generated along t_Q direction, is the field strength for $U(1)_Q$ flux and i is immersion to B . We turned on a flux for $U(1)_Z$ as in (8) thus the corresponding gauge boson acquire mass by Stückelberg mechanism and the symmetry is broken. On the heterotic side, it looks that the anomaly of $U(1)_Z$ is removed by shift of model-dependent axion [36], which is the imaginary part of superfield $T = \int_Q J + i \int_Q B$, where J is the Kähler form, B is the NSNS two-form, and Q is interpreted as two-cycle wrapped by worldsheet instanton [35]. Now T is charged and there is an instanton generating a nonperturbative super potential, guided by $U(1)_Z$ invariance.

To keep $SO(10)$ unification relation for the matter multiplicity, we do not turn on flux along X direction, and the only possible superpotential is of a form $e^{-T} S^n$, $n \in \mathbb{Z}$. In this case, $U(1)_X$ and hypercharge do not belong to the structure group of the vector bundle in the heterotic side, and they may remain as unbroken group in the low energy [36]. Phenomenology of these extra $U(1)$ groups inside E_6 are recently discussed in Ref. [37].

Since we do not turn on flux along X direction, their gauge couplings receive no threshold correction from the flux from F-theory side [3,38]. The four-dimensional gauge coupling is inversely proportional to the volume of four cycle S supporting gauge group, but to be precise it is topologically given by intersection numbers $g_{4D}^{-1} \propto e^{-\phi} \int_S J \wedge J$. Since $SU(3)$ and $SU(2)$ have linearly equivalent cycle [5,39], we have the same four-dimensional coupling. In fact, we have only one gauge coupling of embedded in E_8 , and $SO(10)$, giving the same coupling to $SU(3) \times SU(2) \times U(1)_Y \times U(1)_X$ with the correct normalization in $SO(10)$

$$g_3 = g_2 = \sqrt{\frac{5}{3}} g_Y = \sqrt{40} g_X, \quad \sin^2 \theta_W = \frac{3}{8},$$

at M_s . The $U(1)_X$ can survive as gauge symmetry at relatively low energy scale and would be spontaneously broken down at relative low energy. Threshold corrections for the split Higgs triplets D_1 and \bar{D}_2 would modify the scale.

6. Conclusion

We sought a supersymmetric extension of the Standard Model using spectral cover construction. As a minimal set of conditions, we required the SM group, matter parity, and the correct

number of the Higgs doublet. Each step narrowed the structure group of the vector bundle to a subgroup of $S[U(5) \times U(1)]$, $S[U(4) \times U(1)^2]$, $S[U(3)_\perp \times U(1)^3]$, respectively. Since a smaller structure group such as $S[U(2) \times U(1)^4]$ cannot reproduce the desired spectrum and couplings, the only possible choice in this framework is $S[U(3)_\perp \times U(1)^3]$. Requiring three generations of matter fields and one pair of Higgs doublets, the universal G -flux is turned on with the structure group $S[U(3)_\perp \times U(1)_Z]$, resulting in the multiplicity of the spectrum satisfying $SO(10)$ unification relation. Another flux component along $U(1)_X$ is optional. As a nontrivial consequence of the spectral cover and the resulting matter localization, we are able to distinguish up and down Higgs fields, and obtain a restricted perturbative and nonperturbative superpotentials for the singlets S giving μ -term. Analysis of the consequent dynamics would be an interesting future direction.

Acknowledgements

The author is grateful to Hirotaka Hayashi, Jihn E. Kim, Tatsuo Kobayashi, Bumseok Kyae, Seung-Joo Lee, Hans-Peter Nilles and Martijn Wijnholt for discussions and correspondences. This work is partly supported by the National Research Foundation of Korea (NRF) with grant number 2012-040695.

References

- [1] H. Georgi, S.L. Glashow, Phys. Rev. Lett. 32 (1974) 438; J.C. Pati, A. Salam, Phys. Rev. D 10 (1974) 275.
- [2] H.P. Nilles, hep-th/0410160; J.J. Heckman, C. Vafa, JHEP 0909 (2009) 079.
- [3] R. Donagi, M. Wijnholt, arXiv:0808.2223 [hep-th].
- [4] K.-S. Choi, T. Kobayashi, Phys. Lett. B 693 (2010) 330, arXiv:1003.2126 [hep-th].
- [5] K.-S. Choi, Nucl. Phys. B 842 (2011) 1, arXiv:1007.3843 [hep-th].
- [6] R. Friedman, J. Morgan, E. Witten, Commun. Math. Phys. 187 (1997) 679.
- [7] R. Tatar, T. Watari, Nucl. Phys. B 747 (2006) 212, hep-th/0602238.
- [8] R. Blumenhagen, S. Moster, T. Weigand, Nucl. Phys. B 751 (2006) 186, hep-th/0603015.
- [9] J. Marsano, N. Saulina, S. Schafer-Nameki, JHEP 0908 (2009) 046.
- [10] T.W. Grimm, T. Weigand, arXiv:1006.0226 [hep-th].
- [11] S. Katz, D.R. Morrison, S. Schafer-Nameki, J. Sully, JHEP 1108 (2011) 094, arXiv: 1106.3854 [hep-th].
- [12] K.-S. Choi, H. Hayashi, JHEP 1206 (2012) 009, arXiv:1203.3812 [hep-th].
- [13] R. Donagi, M. Wijnholt, arXiv:0904.1218 [hep-th].
- [14] H. Hayashi, R. Tatar, Y. Toda, T. Watari, M. Yamazaki, Nucl. Phys. B 806 (2009) 224.
- [15] C. Beasley, J.J. Heckman, C. Vafa, JHEP 0901 (2009) 058; C. Beasley, J.J. Heckman, C. Vafa, JHEP 0901 (2009) 059.
- [16] T.W. Grimm, Nucl. Phys. B 845 (2011) 48.
- [17] P. Langacker, J. Wang, Phys. Rev. D 58 (1998) 115010; S.F. King, et al., Phys. Rev. D 73 (2006) 035009.
- [18] See also J.J. Heckman, A. Tavanfar, C. Vafa, arXiv:0906.0581 [hep-th].
- [19] M. Bershadsky, K.A. Intriligator, S. Kachru, D.R. Morrison, V. Sadov, C. Vafa, Nucl. Phys. B 481 (1996) 215; S. Katz, D.R. Morrison, S. Schafer-Nameki, J. Sully, JHEP 1108 (2011) 094.
- [20] R. Blumenhagen, G. Honecker, T. Weigand, JHEP 0506 (2005) 020.
- [21] R. Donagi, Y.H. He, B.A. Ovrut, R. Reinbacher, JHEP 0412 (2004) 054; R. Blumenhagen, S. Moster, R. Reinbacher, T. Weigand, JHEP 0705 (2007) 041.
- [22] K.-S. Choi, B. Kyae, arXiv:1102.0591 [hep-th].
- [23] C.-M. Chen, Y.-C. Chung, JHEP 1103 (2011) 049; C.-M. Chen, Y.-C. Chung, JHEP 1103 (2011) 129.
- [24] R. Tatar, Y. Tsuchiya, T. Watari, Nucl. Phys. B 823 (2009) 1.
- [25] See also C. Ludeling, H.P. Nilles, C.C. Stephan, Phys. Rev. D 83 (2011) 086008; H.M. Lee, et al., Nucl. Phys. B 850 (2011) 1; K.-S. Choi, J.E. Kim, Phys. Rev. D 83 (2011) 065016.
- [26] J. Erler, Nucl. Phys. B 586 (2000) 73, hep-ph/0006051; N. Maekawa, T. Yamashita, Prog. Theor. Phys. 107 (2002) 1201, hep-ph/0202050; M. Cvetic, P. Langacker, in: G.L. Kane (Ed.), Perspectives on Supersymmetry II, World Scientific, 2010, pp. 325–350, hep-ph/9707451.
- [27] For example P.G. Camara, E. Dudas, E. Palti, JHEP 1112 (2011) 112; K.-S. Choi, T. Kobayashi, Nucl. Phys. B 797 (2008) 295; H. Abe, et al., JHEP 0906 (2009) 080.
- [28] U. Ellwanger, C. Hugonie, A.M. Teixeira, Phys. Rept. 496 (2010) 1, and references therein; See also O. Lebedev, S. Ramos-Sánchez, Phys. Lett. B 684 (2010) 48.
- [29] M. Cvetic, J. Halverson, P. Langacker, JHEP 1009 (2010) 076.
- [30] R. Blumenhagen, A. Collinucci, B. Jurke, JHEP 1008 (2010) 079; T.W. Grimm, M. Kerstan, E. Palti, T. Weigand, arXiv:1105.3193 [hep-th]; M. Cvetic, J. Halverson, I. García-Etxebarria, arXiv:1107.2388 [hep-th].
- [31] G.F. Giudice, A. Masiero, Phys. Lett. B 206 (1988) 480; J.E. Kim, H.P. Nilles, Phys. Lett. B 138 (1984) 150.
- [32] P. Fileviez Perez, H. Iminniyaz, G. Rodrigo, Phys. Rev. D 78 (2008) 015013, arXiv:0803.4156 [hep-ph].
- [33] S. Raby, hep-ph/0211024.
- [34] A precise treatment of D -term is made in Ref. [16] and T.W. Grimm, M. Kerstan, E. Palti, T. Weigand, JHEP 1112 (2011) 004, arXiv:1107.3842 [hep-th].
- [35] R. Donagi, M. Wijnholt, arXiv:1005.5391 [hep-th].
- [36] J. Distler, B.R. Greene, Nucl. Phys. B 304 (1988) 1.
- [37] See J. Erler, P. Langacker, S. Munir, E. Rojas, arXiv:1108.0685 [hep-ph], and references therein.
- [38] R. Blumenhagen, Phys. Rev. Lett. 102 (2009) 071601, arXiv:0812.0248 [hep-th].
- [39] K.-S. Choi, JHEP 1002 (2010) 004.