



Hairs of discrete symmetries and gravity



Kang Sin Choi^{a,b}, Jihn E. Kim^{c,d}, Bumseok Kyae^e, Soonkeon Nam^c

^a Scranton Honors Program, Ewha Womans University, Seodaemun-Gu, Seoul 03760, Republic of Korea

^b Center for Fields, Gravity and Strings, CTPU, Institute for Basic Sciences, Yuseong-Gu, Daejeon 34047, Republic of Korea

^c Department of Physics, Kyung Hee University, 26 Gynghedaero, Dongdaemun-Gu, Seoul 02447, Republic of Korea

^d Center for Axion and Precision Physics Research (IBS), 291 Daehakro, Yuseong-Gu, Daejeon 34141, Republic of Korea

^e Department of Physics, Pusan National University, 2 Busandaehakro-63-Gil, Geumjeong-Gu, Busan 46241, Republic of Korea

ARTICLE INFO

Article history:

Received 15 March 2017

Received in revised form 3 April 2017

Accepted 5 April 2017

Available online 13 April 2017

Editor: M. Cvetič

Keywords:

Discrete symmetry

Wormholes

Gravity

Discrete hair

ABSTRACT

Gauge symmetries are known to be respected by gravity because gauge charges carry flux lines, but global charges do not carry flux lines and are not conserved by gravitational interaction. For discrete symmetries, they are spontaneously broken in the Universe, forming domain walls. Since the realization of discrete symmetries in the Universe must involve the vacuum expectation values of Higgs fields, a string-like configuration (hair) at the intersection of domain walls in the Higgs vacua can be realized. Therefore, we argue that discrete charges are also respected by gravity.

© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

It has been known for a long time that discrete sub-groups of gauge groups, the so-called discrete gauge symmetries, are not broken by gravitational interactions [1,2]. Effects of quantum gravity are studied by looking at various topologies of the metric tensor $g_{\mu\nu}$. If some gauge charges are separated from our Universe by metric change, the separated gauge charges cannot be completely hidden from our Universe because they leave long range flux lines. On the other hand, if global charges are separated from our Universe, the lost charges leave no hint to an observer in our Universe and he notices that global charges are not conserved in our Universe. Thus, gauge symmetries are not broken but global symmetries are broken by metric changes. This is the basic reasoning that discrete gauge symmetries are used in particle physics [3]. This top-down approach on discrete symmetries fits to the string compactification [4,5] because string theory does not allow any global symmetry.

In the bottom-up approach, the flux line argument is not so clear. It uses just the classical gauge fields and does not rely on the renormalizability in the theory of elementary particles. To be specific, let us consider a continuous symmetry $U(1)$. If $U(1)$ is a gauge symmetry, it should not have any gauge anomaly. If $U(1)$ is a

global symmetry, it may have a gauge anomaly $U(1)-G-G$ where G is a gauge group as in the Peccei–Quinn (PQ) global symmetry $U(1)_{PQ}$ [6]. Obstructing the PQ symmetry needed for an “invisible” axion was based on this argument [7].

However, the absence of any gauge anomaly is not a guarantee for a gauged $U(1)$ symmetry. Some global $U(1)$ symmetries may not have any gauge anomaly. The difference in the gauge and global symmetries resides in the property on the local transformation, i.e. using a covariant derivative $D_\mu = \partial_\mu - iA_\mu$ in gauge theories, or just an ordinary one ∂_μ in global symmetries. A discrete subgroup of $U(1)$ cannot know whether the mother $U(1)$ is gauged or not. In the bottom-up approach, there must be some other reason for the effects of the metric change.

In this paper, we adopt the concept of “hair” which means that hair’s thickness is the same at any distance from the surface of the head. At the surface, there must be fields at the surface for a hair to be defined. This definition excludes any possibility for hairs of global symmetries. In gauge theories, there are gauge fields at the surface. In gauge theories, the relation of the fields at the surface with the charge Q in the volume enclosed by the surface is provided by the equations of motion and current conservation. Existence of hairs is crucial in guaranteeing the symmetry in the presence of the gravitational interaction. It is known that black holes have gauge-charge hairs, which will be briefly commented in parallel with our method.

E-mail address: jihnekim@gmail.com (J.E. Kim).




Fig. 1. Multiple discrete vacua. Some of minima are shown as green bullets. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

For a gauge charge Q , we have gauge fields spreading out from Q . Consider the current j^μ and the corresponding electric field \mathbf{E} along a line to be interpreted as a hair. We can perform local transformations such that \mathbf{E} is the same along a line but zero outside the line, which behaves as a hair.¹

In this paper, we show how discrete charges can have hairs in the bottom-up approach, and derive that discrete symmetries are not broken by gravity. For an explicit presentation, we will present examples with the Abelian discrete symmetry \mathbf{Z}_N and in particular with \mathbf{Z}_2 illustrations.

2. Discrete charges of \mathbf{Z}_N vacua

A discrete symmetry is defined by the number of minima of the potential V such as in Fig. 1. Let us consider one minimum, say a green bullet in Fig. 1. We can choose the value of the Higgs field to be zero at that point so that the discrete symmetry is realized by the Wigner–Weyl manner. If it has a flat direction there, then one must consider a continuous symmetry, which has been spontaneously broken already. Not considering continuous symmetries, with the multiple vacua of Fig. 1, the discrete symmetry is good at any point of the minima. We will consider the discrete charges at such a minimum.

Realization of discrete symmetries in the Universe leads to domain walls [8]. In the “invisible” axion case [9], the Peccei–Quinn symmetry leads to \mathbf{Z}_N domain walls [10]. For the Kim–Shifman–Vainshtein–Zakharov “invisible” axion where there is only one vacuum [11], even the \mathbf{Z}_1 domain wall can be considered in the Universe evolution [12]. In this case, however, all space points except at the wall are in the same vacuum. Different vacua arise for the cases of $N \geq 2$. Two kinds of walled vacua are possible for \mathbf{Z}_2 , viz. Fig. 2. Two vacua of \mathbf{Z}_2 are defined with discrete charges $q = 2n$ and $2n + 1$, mod. 2 (n = integer).

In Fig. 2(a), the (red, $Q_{\text{total}} = 1$) vacuum is seen from the $q = 0$ (yellow) vacuum. A closed domain wall separates these two. This wall viewed from the yellow vacuum is symbolized by the limegreen color. In Fig. 2(b), the $q = 0$ (yellow, $Q_{\text{total}} = 0$) vac-




Fig. 2. (a) A walled vacuum (red, $q_{\text{total}} = 1$) seen from the $q = 0$ (yellow) vacuum. Inside the wall, the opposite $q = 1$ (red) vacuum is seen through a crack in the wall. This view of the wall is colored limegreen. (b) A walled vacuum (yellow, $q_{\text{total}} = 0$) seen from the $q = 1$ (red) vacuum. This view of the wall is colored blue. Dashes represent closing surfaces and black dots represent particles. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

uum is seen from the $q = 1$ (red) vacuum. The wall viewed from the red vacuum is symbolized by the blue color. In Fig. 2(a), the dashed boundary encloses the walled $q = 1$ vacuum. A scalar field ϕ in the $q = 0$ or $q = 1$ vacua is represented by $e^{iq\pi} R(\mathbf{x})$.

Let us illustrate examples in \mathbf{Z}_2 . Then, q can be 0 or 1. For $q = 0$, we use the field VEV $\phi = 0$.² For a ball of discrete charge q , the radius of the ball is determined by minimizing the energy

$$E_\omega = E + \omega \left[q - \frac{1}{2i} \int d^3x (\Phi^* \partial_t \Phi - \Phi \partial_t \Phi^*) \right], \quad (1)$$

where ω (with the energy dimension) is the Lagrange multiplier and q of the ball can be defined as

$$q = \frac{1}{2i} \int d^3x (\Phi^* \partial_t \Phi - \Phi \partial_t \Phi^*). \quad (2)$$

In the evolving Universe, the vacuum inside the ball expands such that ϕ in the red becomes constant. For a spherically symmetric $R(\mathbf{x})$, let us parametrize it as

$$\Phi = \sqrt{\frac{3k^3}{4\pi^4\omega}} e^{i\omega t} \begin{cases} 1, & \text{for } 0 \leq r < \frac{\pi}{k} \\ 0, & \text{for } r > \frac{\pi}{k} \end{cases} \quad (3)$$

where π/k is the radius of the ball, and $\Phi = 0$ in the yellow part of Fig. 2(a). So, we obtain

$$\frac{1}{2i} \int_{(\text{inside dashed})} d^3x (\Phi^* \partial_t \Phi - \Phi \partial_t \Phi^*) = 1.$$

The total charge q inside the dashed surface of Fig. 2(a) is 1, and the dashed string symbolizes this fact.

Definition of charge q by Eq. (2) is not by the Nöther current. It is simply defined by the vacuum expectation value (VEV) of the phase of a Higgs field Φ . To relate this charge q to the charge defined by the Nöther current, the t dependence of Φ is introduced as the example in Eq. (3). To make it an integer, the VEV which is designed as a constant³ is appropriately chosen. Equation (1) is the matching condition to the charge q calculated by the Nöther current. Discrete symmetries in the Universe are realized by the VEVs of Higgs field Φ having degenerate minima as shown in Fig. 1. So, it is appropriate to figure out the discrete charges in the vacuum

¹ Here, the line is not a mathematical one but has some physical thickness. Thus, gauge charges can have hairs but global charges cannot, and metric changes know only hairs.

² If $\phi = -v$ corresponds to $q = 0$, we add a constant v to simplify the value of ϕ .

³ In the connected portion in the Universe, the minimum of the potential with a fixed value of Φ is chosen everywhere.




Fig. 3. (a) A \mathbf{Z}_N walled ball seen in the $q=0$ vacuum. The tail of the “tadpole” can have $n=0, 1, \dots, N-1$ dashed lines. N dashed line is equivalent to no tail, i.e. no \mathbf{Z}_N charge. (b) Another view of (a) by the discrete flux, and (c) an expanded view of (b) with the dashed boundary touching the wall. In (a), the discrete charges inside two dashed surfaces are the same.

of spin-0 bosons as shown in Fig. 2 where the limegreen surface separates two different Higgs portions in the Universe. In case of Fig. 2(b), we apply discrete transformation $e^{i\pi Q}$. Then it is equivalent to the discussion in Fig. 2(a).

In Fig. 2(b), $q_{\text{total}} = 4 = \text{even} = 0$ within the white dashed surface, which must be the case for any closed surface. So, the white dashed surface of Fig. 2(b) can be shrunk to naught. In Fig. 2(a), $q_{\text{total}} = 3 = \text{odd} = 1$ within the dashed surface, which must be the case for any closed surface, which encloses the limegreen ball. To identify q_{total} , we extend a dashed line from the limegreen ball up to the horizon. The dashed surface of Fig. 2(a) is named as an infinite-tail “tadpole”, having a head and a long tail. At the limegreen surface, there exists the Higgs field which provides the logistics for a hair to exist in our case.

Suppose that the U(1) charges of Φ and ψ are 2 and 1, respectively, and there is no U(1)–G–G anomaly where G is a gauge group. Then, this U(1) can be a gauge or global group. The U(1) transformations by an angle θ on the fields are $\Phi \rightarrow e^{i4\pi\theta}\Phi$ and $\psi \rightarrow e^{i2\pi\theta}\psi$. A potential invariant under U(1) is

$$V = -\left(\sqrt{2}\mu\psi^{*2}\Phi + \text{h.c.}\right) + \dots \quad (4)$$

where \dots are other U(1) invariant terms. At a high scale f , the U(1) is broken to \mathbf{Z}_2 by the VEV, $\langle\Phi\rangle = f/\sqrt{2}$. Let us split ψ to a radial and phase fields,

$$\psi = \frac{v+\rho}{\sqrt{2}} e^{i2\pi\theta} \quad (5)$$

For a moment, let $\langle v \rangle = 0$ where $\theta = \phi/v$. At this stage, there is no θ dependence in the potential because $\langle\phi\rangle = 0$. There is no way to distinguish the U(1) as a global or gauge group. Now, let $|\psi|$ develop a VEV, $\langle|\psi|\rangle = v/\sqrt{2}$, so that the \mathbf{Z}_2 symmetric potential, $-(f\mu\psi^{*2} + \text{h.c.})$, develops a θ dependence,

$$V = -f\mu v^2 \cos\theta + \dots \quad (6)$$

In the region $\phi = -v$ (yellow) and $+v$ (red), let $q=0$ (yellow) and 1 (red), respectively.⁴ Now, the difference appears. In a gauged U(1), θ becomes redundant in the sense that it can be removed from V by a gauge transformation [2]. But, a global U(1) manifests itself in the form of the domain-wall energy density shown in Eq. (6). The effective thickness of the domain wall is

$$\lambda_{\text{thickness}} \approx \frac{\sqrt{\lambda}}{v} \sqrt{\frac{f}{\mu}} \quad (7)$$

⁴ The discrete charges are the phase values of scalar fields.

where λ is a quartic coupling constant, determining f from the neglected terms in (4). So, the domain wall can be seen as we discussed above, and the intersection of domain walls works as a hair.

3. Discrete charge conservation in the bottom-up approach

A conserved current of a continuous symmetry is constructed with an infinitesimal shift of fields, $\Psi \rightarrow e^{i2\pi\epsilon(x)Q}\Psi$ where Q is the generator of the transformation. For a discrete charge I of Ψ in \mathbf{Z}_N symmetry, we adopt this around a specific discrete value I with $I = \{1, 2, \dots, N-1\}$,

$$\Psi \rightarrow e^{i2\pi\left[\frac{I}{N} + \epsilon(x)\right]}\Psi. \quad (8)$$

The vacua of fields are defined for specific values of scalar fields as in Sec. 2, which is related to the charges of quanta, Eq. (8), in the volume via conservation of the current.

For a discrete symmetry, the vacuum structure of Higgs fields allows a hair(s) as shown in Sec. 2. This string-like hair starts from a nonzero discrete charge q and ends at another discrete charge or at the horizon. In Fig. 3(a), we show \mathbf{Z}_N hairs spreading out from a nonzero discrete charge. The Higgs vacua are defined by the VEVs of Higgs fields. In Fig. 3(a), we enclose the discrete charges by two dashed surfaces, the thick and thin ones. The discrete charges inside the thin and thick surfaces are exactly the same. If we assign one discrete charge to one dashed line, the discrete charges inside the thin and thick surfaces are exactly the same, i.e. the hair interpretation of the dashed lines equates these two estimations. For example, the Higgs vacuum value at the star at the dashed line is the same as that at the triangle. Calculating the discrete charge in Fig. 3(b) is like calculating the discrete charge in Fig. 3(c) where we made it clear by moving the dashed surface touching the wall at the RHS of Fig. 3(c). So, when discrete charges move, we can consider them dragging dash lines corresponding to some units of discrete charges.

Now let us proceed to discuss the wormhole effects in the metric theory of gravity. Through wormholes, discrete charges can flow out from our Universe O. For simplicity, now let us focus on \mathbf{Z}_2 . An infinite-tail “tadpole” is symbolized in Fig. 4(a), where $Q_{\text{total}} = 1$. Discrete charge flow can be visualized as a “tadpole” passing through the wormhole as shown in Fig. 4(b). If one tries to separate out the shadow world S from O by cutting the wormhole through the gray plane in Fig. 4(b), it cuts the infinite tail of “tadpole”. Then, at the cut plane, dashed lines are attached to a walled ball at each surface as shown in Fig. 2(c). Recovering the wormhole throat, the observer O confirms that no discrete charge is lost. To the observer O, gravitational effects do not break the dis-




Fig. 4. (a) A walled ball seen in the $q=0$ vacuum. The tail of a tadpole-like configuration extends to the horizon. (b) A tadpole passing through a wormhole. (c) The wormhole cut in the gray plane in (b). The observer O recovers the discrete charge. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

crete symmetry in consideration. This type of wormhole argument was used for a U(1) gauge symmetry in Ref. [5].

4. Scalar vacua

Let us illustrate how such hairs from discrete charges are set up in the spontaneously broken \mathbf{Z}_N vacua. For simplicity, we present it in \mathbf{Z}_2 . Let us consider a dashed surface of Fig. 2(a). Our objective is to obtain a two dimensional delta function at the surface of radius r in the spherical-polar coordinate system, (θ_0, φ_0) . Consider an effective \mathbf{Z}_2 symmetric action,

$$\mathcal{L}_{\text{eff}} = \frac{1}{M_{\text{eff}}^2} (\partial^\mu \partial^\nu \psi^*) (\partial_\mu \partial_\nu \psi). \quad (9)$$

At a closed surface, we can consider an effective Lagrangian where the Lorentz symmetry is broken as in Fig. 2(a), and contract with the (μi) indices in (9), $j^{\mu i}$. To have currents related to (9), consider a shift in the discrete vacuum I , viz. Eq. (8). There are two index current, proportional to $\partial_\mu \partial_i \varepsilon(x)$, and one index currents, proportional to $\partial_\mu \varepsilon(x)$ and $\partial_i \varepsilon(x)$. One index currents are not of interest here. The two-index current is

$$\propto i e^{i 2\pi (\frac{I}{N} - \frac{I}{N})} (\psi^* [Q_\psi \partial^\mu \partial^i \psi] - [Q_\psi \partial^\mu \partial^i \psi]^* \psi). \quad (10)$$

Let us consider the two-index phase at $I=1$ and $N=2$, i.e. for $Q_\psi = 1$ and $\psi = \frac{f+\rho}{\sqrt{2}f} e^{i 2\pi \phi/f}$,

$$\begin{aligned} \frac{1}{2\pi i} j^{ij} &= \frac{f}{M_{\text{eff}}^2} \partial^i \partial^j \phi \rightarrow Q = \frac{f}{M_{\text{eff}}^2} \int d^3 x \phi \partial^i \partial^j \phi \\ &= \frac{f \lambda_{\text{thickness}}}{M_{\text{eff}}^2} \int d^2 x \partial^i \partial^j \phi \\ &= \sqrt{\frac{\lambda v}{\mu}} \int d\Omega \frac{f v}{M_{\text{eff}}^2} j^{\theta\varphi}, \end{aligned} \quad (11)$$




Fig. 5. A \mathbf{Z}_2 string.

where v is the VEV of ϕ breaking \mathbf{Z}_2 , μ is the ϕ mass as the result of this breaking, and $\lambda_{\text{thickness}}$ is given in Eq. (7). A delta function is obtained from the derivative of a step function in the angle direction. It is shown in Fig. 5. Here, the parameter M_{eff} is defined at the radius r so that Q becomes 1,

$$M_{\text{eff}}^2 = \sqrt{\lambda \frac{f^2 v^3}{\mu}}, \quad (12)$$

and

$$j^{\theta\varphi}(\theta, \varphi) = \frac{1}{r^2} \delta(\cos \theta - \cos \theta_0) \delta(\varphi - \varphi_0). \quad (13)$$

With the current conservation, this surface integral is related to the head charge of the tadpole,

$$\partial_\mu j^{\mu i} = 0 \rightarrow Q^i \propto \int_V d^3 x \frac{\partial}{\partial x^j} j^{ji} = \int_{\Sigma_r} d^2 \sigma_r j^{ri}, \quad (14)$$

where Σ_r is the surface orthogonal to the radial direction. The current conservation used in (14) is obtained via the equation of motion $\partial^2 \psi = -m^2 \psi$ and $\partial^2 \psi^* = -m^2 \psi^*$,

$$\begin{aligned} \partial_\mu j^{\mu i} &= \frac{1}{2} [\psi^* \partial^i \partial^2 \psi + (\partial_\mu \psi^*) \partial^i \partial^\mu \psi \\ &\quad - (\partial^2 \psi^*) \partial^i \psi - (\partial_\mu \psi^*) \partial^i \partial^\mu \psi] \\ &= \frac{1}{2} [\psi^* \partial^i \partial^2 \psi - (\partial^2 \psi^*) \partial^i \psi] = 0. \end{aligned} \quad (15)$$

5. Conclusion

Showing that discrete charges carry hairs in the bottom-up approach, we argued that discrete symmetries are respected by gravity. It is based on the fact that discrete symmetries are realized with domain walls in the Universe via the VEVs of Higgs fields, and the intersection of domain walls looks like a hair.

Acknowledgements

K.S.C. is supported in part by NRF-2015R1D1A1A01059940, J.E.K. is supported in part by the National Research Foundation (NRF) grant funded by the Korean Government (MEST) (NRF-2015R1D1A1A01058449) and the IBS (IBS-R017-D1-2016-a00), and B.K. is supported in part by the NRF-2013R1A1A2006904.




Fig. 6. A charged black hole.




Fig. 7. A field vacua around a black hole.

Appendix A. Around a blackhole

Around a blackhole, the argument goes in parallel to the case of wormhole. So, we briefly point out the flux line argument in a charged Universe and draw the similarity in case of discrete charges.

The larger event horizon (out of two solutions) of the Reissner-Nordström black hole [13] occurs at

$$r_+ = \frac{1}{2} \left(r_s + \sqrt{r_s^2 - 4r_Q^2} \right) \quad (16)$$

where r_s is the Schwarzschild radius without the charge. In Fig. 6(a), we show the situation. The event horizon takes into account the energy inside it, as depicted with the white illustration. It is basically the impossibility of the graviton field to go out of the horizon, bounded by $T_{\mu\nu}$. Any graviton, a spin-2 field, can end at another energy point, due to the positive energy theorem, inside the blackhole horizon. When we consider mass of a charged particle, it includes the field energy also. This count of energy subtracts the electromagnetic field energy permeating from the horizon to infinity, which is shown as r_Q^2 in Eq. (16). Within the closed boundary shown in Fig. 6(a), the electromagnetic field cannot end inside the blackhole horizon for a net charge Q given inside the black hole. So, the field line goes out of the horizon of the gravity field, which looks like a hair from the black hole. Of course, the outside field is not included in $T_{\mu\nu}$ on the right-hand side of the Einstein equation. So, if fields permeating over the whole space are present, it is better to consider their effects just inside the blackhole. In Fig. 7 around a closed surface black hole, the electromagnetic field configuration, not interfering with the fact of the closed surface nature of the black hole, is illustrated.

Now, let us consider that the boundary of the fields is confined inside the horizon. If the field lines are forbidden to cross





Fig. 8. Hairs of a black hole: (a) for an electromagnetic charge, and (b) for a discrete tadpole.

Fig. 9. A tadpole is thrown into the $q = 0$ vacuum.

the boundary for the spin-1 fields also, one cannot allow a net charge Q inside the horizon and must give up the closed surface of the blackhole,⁵ and the metric must allow the open geometry as shown in Fig. 6(b). By gauge transformation, the electromagnetic fields of Fig. 6(a) are transformed to those of Fig. 6(b) where, in the most part of the blackhole space, the gravity field is bounded within the dashed area but the electromagnetic field goes out into the open space through the pinched hole. This is the hair we explain within our set-up. Spin-0 fields do not have the flux lines but can have different quantum numbers distinguishing different domains of the spin-0 field vacua. For spin-1 field the hair is the field strength and for spin-0 field we argued that it must be the intersection of domain walls. This was named as 'tadpole'. In Fig. 8, the hairs of U(1) gauge field and Z_2 tadpole are shown. In Fig. 8(b), the discrete vacuum of the blackhole in the open geometry is $q = 0$. Here, we note the difference between the flux line and the discrete tadpole. The electromagnetic flux carries energy and hence affects the blackhole radius. But, the discrete vacua in Fig. 1 are degenerate and different vacua have the same energy. If they have strings (intersection of domain walls), they must be taken into account in the energy calculation inside the blackhole, not considering the outside part. So, the blackhole radius is not changed.

Vacua of spin-0 field are not constrained by the blackhole horizon, and the same vacuum can be connected to the outside as the field lines of Fig. 6(a). In Fig. 9, a Z_2 tadpole is thrown into the $q = 0$ blackhole vacuum, and the outside observer notices that he lost a discrete charge $q = 1$ to the blackhole. Namely, he notices that the black hole has a Z_2 hair. If the tail is cut, the outside observer notices that the blackhole ate only even number of Z_2 charge with some energy increase inside the blackhole but the discrete charge is not increased in the blackhole. Thus, the existence of a blackhole does not violate the discrete symmetry.

⁵ In the closed Universe, a non-zero charge must break the U(1) gauge symmetry [14].

References

- [1] L.M. Krauss, F. Wilczek, Discrete gauge symmetry in continuum theories, *Phys. Rev. Lett.* 62 (1989) 1221, <http://dx.doi.org/10.1103/PhysRevLett.62.1221>.
- [2] T. Banks, Effective Lagrangian description of discrete gauge symmetries, *Nucl. Phys. B* 323 (1989) 90, [http://dx.doi.org/10.1016/0550-3213\(89\)90589-0](http://dx.doi.org/10.1016/0550-3213(89)90589-0).
- [3] L.E. Ibañez, G.G. Ross, Discrete gauge symmetries and the origin of baryon and lepton number conservation in supersymmetric versions of the standard model, *Nucl. Phys. B* 368 (1992) 3, [http://dx.doi.org/10.1016/0550-3213\(92\)90195-H](http://dx.doi.org/10.1016/0550-3213(92)90195-H).
- [4] T. Kobayashi, H.P. Nilles, F. Plöger, S. Raby, M. Ratz, Stringy origin of non-Abelian discrete flavor symmetries, *Nucl. Phys. B* 768 (2007) 135, [arXiv:hep-ph/0611020](http://arxiv.org/abs/hep-ph/0611020).
- [5] J.E. Kim, Abelian discrete symmetries Z_N and Z_{nR} from string orbifolds, *Phys. Lett. B* 726 (2013) 450, [arXiv:1308.0344 \[hep-th\]](http://arxiv.org/abs/1308.0344).
- [6] R.D. Peccei, H.R. Quinn, CP conservation in the presence of instantons, *Phys. Rev. Lett.* 38 (1977) 1440, <http://dx.doi.org/10.1103/PhysRevLett.38.1440>.
- [7] S.M. Barr, D. Seckel, Planck-scale corrections to axion models, *Phys. Rev. D* 46 (1992) 539, <http://dx.doi.org/10.1103/PhysRevD.46.539>; M. Kamionkowski, J. March-Russel, Planck-scale physics and the Peccei–Quinn mechanism, *Phys. Lett. B* 282 (1992) 137, [arXiv:hep-th/9202003](http://arxiv.org/abs/hep-th/9202003); R. Holman, S.D.H. Hsu, T.W. Kephart, R.W. Kolb, R. Watkins, L.M. Widrow, Solutions to the strong CP problem in a world with gravity, *Phys. Lett. B* 282 (1992) 132, [arXiv:hep-ph/9203206](http://arxiv.org/abs/hep-ph/9203206).
- [8] Ya.B. Zeldovich, I.Yu. Kobzarev, L.B. Okun, Cosmological consequences of the spontaneous breakdown of discrete symmetry, *Zh. Eksp. Teor. Fiz.* 67 (1974) 3–11, *JETP* 40 (1974) 1–5.
- [9] J.E. Kim, Light pseudoscalars, particle physics and cosmology, *Phys. Rep.* 150 (1987) 1, [http://dx.doi.org/10.1016/0370-1573\(87\)90017-2](http://dx.doi.org/10.1016/0370-1573(87)90017-2).
- [10] P. Sikivie, Of axions, domain walls and the early Universe, *Phys. Rev. Lett.* 48 (1982) 1156, <http://dx.doi.org/10.1103/PhysRevLett.48.1156>.
- [11] J.E. Kim, Weak interaction singlet and strong CP invariance, *Phys. Rev. Lett.* 43 (1979) 103, <http://dx.doi.org/10.1103/PhysRevLett.43.103>; M.A. Shifman, V.I. Vainshtein, V.I. Zakharov, Can confinement ensure natural CP invariance of strong interactions?, *Nucl. Phys. B* 166 (1980) 493, [http://dx.doi.org/10.1016/0550-3213\(80\)90209-6](http://dx.doi.org/10.1016/0550-3213(80)90209-6).
- [12] A. Vilenkin, A.E. Everett, Cosmic strings and domain walls in models with Goldstone and pseudo-Goldstone bosons, *Phys. Rev. Lett.* 48 (1982) 1867, <http://dx.doi.org/10.1103/PhysRevLett.48.1867>; S.M. Barr, K. Choi, J.E. Kim, Axion cosmology in superstring models, *Nucl. Phys. B* 283 (1987) 591, [http://dx.doi.org/10.1016/0550-3213\(87\)90288-4](http://dx.doi.org/10.1016/0550-3213(87)90288-4).
- [13] H. Reissner, Über die Eigengravitation des elektrischen Feldes nach der Einsteinschen Theorie, *Ann. Phys.* 50 (1916) 106; G. Nordström, On the energy of the gravitational field in Einstein's theory, *Verhandl. Koninkl. Ned. Akad. Wetenschap., Afdel. Natuurk., Amsterdam* 26 (1918) 1201.
- [14] J.E. Kim, T. Lee, U(1) gauge symmetry breaking in a charged closed Universe, *Mod. Phys. Lett. A* 5 (1990) 2209, <http://dx.doi.org/10.1142/S0217732390002511>.