



A Photolabile Backbone Amide Linker for the Solid-Phase Synthesis of Cyclic Peptides and C-terminal Thioesters

Kaif R. Khan¹ · Mary L. Niedrauer² · Matthew A. Hostetler³ · Soo Sung Kang⁴ · Mark A. Lipton⁵

Received: 19 July 2025 / Accepted: 11 August 2025 / Published online: 8 September 2025
© The Author(s) 2025

Abstract

Introduction A new backbone amide linker has been developed for the synthesis of cyclic and C-terminally modified peptides that permits photochemical detachment of the synthesized peptide from the solid support, thus avoiding problems associated with acid deprotection conditions.

Methods An initial survey of known photolabile motifs for their ability to produce a linker-bound model dipeptide in high yield and their ability to undergo efficient photochemical detachment of the model dipeptide found that the 6-nitroveratryl (Nve) motif afforded the most efficient release of the dipeptide. The problematic acylation of Nve-bound amino esters was solved through the development of the 2-hydroxy-4-carboxy-6-nitrobenzyl (Hcnb) linker, which utilizes an O-to-N transacylation to afford efficient acylation of even sterically hindered, linker-bound amino esters. Two different approaches were carried out to avoid/minimize diketopiperazine formation. This methodology was then used to synthesize cyclic peptides and peptide thioesters.

Results The Hcnb linker was found to afford high yields of amino acid loading, acylation, and photolytic cleavage of model tripeptides. Attachment of the Hcnb linker to the aminomethyl TG resin permitted the solid phase synthesis of representative cyclic peptides and C-terminal thioesters in high overall yield and purity.

Conclusion Hcna is a new photolabile backbone amide linker that has been used to synthesize cyclic peptides and peptide thioesters. It was found that the linker efficiently released the synthesized peptides when UV light was shed on it. This linker is also stable to acids meaning global deprotections can be done on resin and can be washed to remove by-products, releasing only the peptide upon its photocleavage from the resin.

Keywords Peptide synthesis · Photochemistry · Peptide cyclization · Thioester formation

Introduction

The head-to-tail cyclization of peptides (Fusetani 1993; Humphrey 1997; Katsara 2006) is a common and important strategy, employed both in nature and in synthetic applications, for improving the bioavailability and conformational rigidity of polypeptides (Kessler 1982; Samanen 1991; Biron 2008). The lack of either an N- or C-terminus increases proteolytic stability while internal hydrogen bonding and a more compact structure improve the membrane permeability of cyclic peptides (White 2011; Dougherty 2019). These advantages are reflected in the prominence of cyclic peptides among peptidic therapeutic agents (Veber 1981; Darkin-Rattray 1996; Coulouarn 1998; Foister 2006), such as gramicidin S (Gause 1944) and cyclosporine A (Shevach 1985). The chemical synthesis of cyclic peptides, however, has been complicated by the traditional approach

✉ Mark A. Lipton
lipton@purdue.edu

¹ Present address: Abbott Laboratories, 100 Abbott Park Road, Abbott Park, IL 60064, USA

² Present address: Low Institute for Therapeutic Research, West Lafayette, IN 47906, USA

³ Present address: Evonik Industries, 1650 Lilly Rd, Lafayette, IN 47909, USA

⁴ Present address: Department of Pharmacy, Ewha Womans University, 52, Ewhayeodae-gil, Seodaemun-gu, Seoul 03760, Republic of Korea

⁵ Department of Chemistry, Purdue University, 560 Oval Drive, West Lafayette, IN 47907, USA

to solid-phase peptide synthesis using the C-terminal carboxylate to covalently attach the growing peptide to a polymeric support. The most straightforward approach involving detachment of a fully protected linear peptide precursor from the resin followed by macrocyclization in solution is complicated by solubility problems, oligomerization and C-terminal epimerization reactions. To avoid the problems associated with head-to-tail cyclization in solution, various approaches have been devised for on-resin cyclization of peptides, including Kaiser's Oxime resin (Mihara 1995), Richter's thioester linker (Richter 1994), Waldmann's hydrazide linker (Rosenbaum 2001), Kenner's sulfonamide linker (Kenner 1971) and side-chain attachment of the linear peptide to the resin (Krishnamoorthy 2006). An important general strategy for the on-resin, head-to-tail cyclization of peptides was introduced by the "backbone amide linker" (BAL) strategy of Albericio and Barany (Jensen 1998). In this approach, the nitrogen atom of an internal amide bond is used to tether the growing peptide to the polymeric resin, thereby liberating both N- and C-termini for modification. It is noteworthy that this strategy also addresses the more general problem of C-terminal modification of the resin-bound peptide. Since its introduction, the BAL strategy has been employed by a number of researchers and several modifications have been developed (Bourne 1999; Bourne 2001; Boas 2002; Brask 2003; Boas 2004; Pittelkow 2005; Jessing 2006; Liley 2006; Pittelkow 2006). Additional applications of the BAL strategy involve the synthesis of C-terminal modifications, such as trithioorthoesters (Guillaumie 2000; Kappel 2005), aldehydes (Mori 2004; Soural 2007; Springer 2008), or functionalized amides.

Another application of the C-terminal modification of peptides was revealed by the discovery of the native chemical ligation of peptide bonds by Kent, Muir and Dawson (Dawson 1994). Their use of peptide thioesters to couple with N-terminal cysteine residues has necessitated the development of various methods for the solid-phase synthesis of peptide thioesters (Mende 2011; Li and Dong 2017). Since that landmark discovery, related processes such as Staudinger ligation (Nilsson 2000) have been developed that likewise depend on the availability of peptide thioesters. In response to this need, a number of creative approaches to the solid-phase synthesis of peptide thioesters have been developed, yet the availability of C-terminal modification in

the BAL strategy remains one of the most straightforward approaches to the synthesis of peptide thioesters.

Although acidic deprotection conditions are widely used in peptide synthesis, there are situations involving acid-sensitive functionality that may require alternative deprotection conditions. For these cases, other methodologies have been explored and developed (Lloyd 1993; Guillier 2000; Kessler 2003). One of the most appealing ideas has been to use photochemical detachment from the resin as an extra dimension of protecting group orthogonality in solid phase peptide synthesis. Photolabile linkers (Gallop 1994; Heidler 2005; Kang 2008; Liang 2016; McAllister 2005; Scott 2006), in which long-wave UV radiation (300–360 nm) is used to detach an organic functional group – usually a carboxylic acid, a carbamate (a latent amine), an alcohol, or a phosphate – from a polymer or glass surface, provide an alternative to the classic acidic or basic cleavage protocols used in solid-phase organic synthesis. Ideally, photochemical deprotection should proceed with high functional group selectivity, low amounts of side reactions and involve no separation of product from reactants. Photochemical deprotection has also found favor with applications in the parallel synthesis of compound libraries (Sheehan 1973).

Herein we describe studies directed toward the development of a photolabile BAL linker (Fig. 1) to address the problems associated with the BAL strategy as a general method for the synthesis of cyclic and C-terminally modified peptides. Our studies began with a survey of several known photolabile motifs for their applicability to BAL linker design (Fig. 1). The candidate photolabile motifs studied were the *p*-oxyphenacyl (Op) (Bellof 1985), α -methyl-*p*-oxyphenacyl (Mop) (Sheehan 1964; Peach 1995), *p*-oxybenzoin (Obz) (Bergmark 1985), α,α -dimethyl phenacyl (Dmp) (Patchornick 1970) and 6-nitroveratryl (Nve). All of these photolabile groups undergo facile n to π^* or π to π^* excitation at 300–360 nm and release oxygen-based substituents with few side reactions. Appropriate synthetic precursors 1–5 were chosen for each of the motifs, as shown in Fig. 2.

Consequently, our preliminary survey focused on the Op, Mop, Obz, Dmp, and Nve motifs, their preparation and photolysis. Of specific interest were (a) the ease of anchoring amino acids to the linker, (b) the rate and efficiency of photocleavage, and (c) the suppression of side reactions





Fig. 1 General concept of amino acid anchoring and photolytic deprotection

Fig. 2 Candidate photolabile backbone amide linkers and their synthetic precursors



during photolytic deprotection. To assess these questions, the dipeptide Boc-Phe-Gly-OMe (**8**) was chosen as a model system. After the photolabile motif was chosen, effort was directed toward its attachment to a solid support compatible with the solvents and reagents common to peptide synthesis. Finally, the problems of diketopiperazine formation, photocleavage optimization, and maximization of cyclization yields were addressed.

Materials and methods

Synthesis of Dmp

1-(2,5-dimethyl-4-alkylphenyl)-propanone (4b) To a mixture of propionyl chloride (0.51 mL, 5.88 mmol) and AlCl₃ (780 mg, 5.88 mmol) in CH₂Cl₂ (25 mL) was added a solution of 4-(2,5-dimethylphenyl)butanoic acid (**4a**, 377 mg, 1.96 mmol) in CH₂Cl₂ (5 mL) at 0 °C. The reaction mixture was stirred for 1 h at the same temperature and poured into cold HCl solution (40 g ice and 10 mL of conc. HCl). The organic material was extracted by addition of CH₂Cl₂ (25 mL) and the aqueous layer was extracted with another portion of CH₂Cl₂ (50 mL). The combined organic layer was washed with brine (50 mL), dried with anhydrous MgSO₄, passed through 3 cm silica gel pad, and evaporated under reduced pressure. The crude product was dissolved in oxalyl chloride (3 mL) and stirred overnight. After removal extra

oxalyl chloride by evaporation under reduced pressure, the organic material was dissolved in CH₂Cl₂ (30 mL). To this solution, benzyl amine (0.257 mL, 2.35 mmol) and DIEA (1.7 mL, 9.8 mmol) were added and stirred 3 h. The organic layer was washed with brine (30 mL), dried with anhydrous MgSO₄, evaporated under reduced pressure and purified by flash column chromatography using 33% EtOAc in Hexanes to give **4b** as a pale-yellow solid.

1-(2,5-dimethyl-4-alkylphenyl)-1-hydroxypropan-2-one oxime (4c) To a solution of **4b** (0.51 g, 1.50 mmol) and isoamyl nitrite (0.24 mL, 1.80 mmol) in MeOH (10 mL) at 45 °C was added dropwise to concentrated HCl (0.25 mL) and the resulting mixture was stirred for 3 h at the same temperature. After cooling to room temperature, the organic material was partitioned into EtOAc (40 mL) and H₂O (40 mL). The organic layer was washed with brine (40 mL), dried with anhydrous MgSO₄, passed through 3 cm silica gel pad, and evaporated under reduced pressure. The residue was dissolved in MeOH (15 mL) and NaBH₄ (50 mg) was added under ice bath. After stirring for 30 min, the organic material was partitioned into EtOAc (40 mL) and HCl solution (0.1 M, 30 mL). The organic layer was washed with saturated NaHCO₃ (30 mL) and with brine (30 mL), dried over anhydrous MgSO₄, and evaporated under reduced pressure. The residue was purified by flash column chromatography using 66% EtOAc in hexanes to give **4c** as a white solid.

1-(2,5-dimethyl-4-alkylphenyl)-1-hydroxypropan-2-one (4) To a solution of **4c** (0.30 g, 0.81 mmol) in DMF (2

mL) was added glyoxylic acid (60% water solution, 5 mL). After stirring for 5 h, the organic material was partitioned into EtOAc (40 mL) and H₂O (40 mL). The organic layer was washed with brine (2×30 mL), dried with anhydrous MgSO₄, evaporated under reduced pressure, and the residue was purified by flash column chromatography using 50% EtOAc in hexanes to give **4** as a pale-yellow oil.

Alkylation of Amino Ester with Photolabile Linker 1–5 (**6a–e**)

H-[Op]-Gly-OMe (6a). To a solution of **1**, (150 mg, 0.66 mmol) in anhydrous CH₃CN (5 mL) were added GlyOMe·HCl (248 mg, 1.98 mmol), and DIEA (0.52 mL, 2.97 mmol). The reaction mixture was stirred overnight at room temperature, evaporated under reduced pressure and partitioned into CH₂Cl₂ (20 mL) and H₂O (20 mL). The organic layer was washed with brine (20 mL), dried with anhydrous MgSO₄, evaporated under reduced pressure, and purified by flash column chromatography using 33% EtOAc in hexanes to give **6a** as a pale-yellow oil.

H-[Mop]-Gly-OMe (6b). To a solution of **2**, (200 mg, 0.63 mmol) in anhydrous CH₃CN (8 mL) were added GlyOMe·HCl (237 mg, 1.89 mmol), DIEA (0.49 mL, 2.84 mmol) and tetrabutylammonium iodide (25 mg). The reaction mixture was refluxed for overnight, cooled to room temperature, evaporated under reduced pressure, and partitioned into CH₂Cl₂ (20 mL) and H₂O (20 mL). The organic layer was washed with brine (20 mL), dried with anhydrous MgSO₄, evaporated under reduced pressure, and purified by flash column chromatography using 33% EtOAc in hexanes to give **6b** as a colorless oil.

H-[Obz]-Gly-OMe (6c). To a solution of **3**, (123 mg, 0.39 mmol) in anhydrous MeOH (20 mL) was added GlyOMe·HCl (242 mg, 1.93 mmol). The reaction mixture was refluxed for 24 h, cooled to room temperature, evaporated under reduced pressure, and partitioned into CH₂Cl₂ (20 mL) and H₂O (20 mL). The organic layer was washed with brine (20 mL), dried with anhydrous MgSO₄, evaporated under reduced pressure, and purified by flash column chromatography using 33% EtOAc in hexanes to give **6c** as a pale-yellow oil.

H-[Dmp]-Gly-OMe (6d). To a solution of **4**, (200 mg, 0.57 mmol) in anhydrous toluene (20 mL) was added GlyOMe·HCl (215 mg, 1.71 mmol). The reaction mixture was refluxed for 5 h, cooled to room temperature, and partitioned into EtOAc (20 mL) and H₂O (40 mL). The organic layer was washed with brine (40 mL), dried with anhydrous MgSO₄, evaporated under reduced pressure, and purified by flash column chromatography using 33% EtOAc in hexanes to give **6d** as a colorless oil.

H-[Nve]-Gly-OMe (6e). To a solution of **5** (515 mg, 2.44 mmol) in MeOH (20 mL), were added GlyOMe·HCl (459 mg, 3.66 mmol), sodium bicarbonate (348 mg, 4.15 mmol). The reaction mixture was stirred for 4 h, cooled to 0 °C. After portionwise addition of NaBH₄ (6.0 mmol, 227 mg), the mixture was stirred for 30 min and poured into HCl solution (0.1 N, 50 mL). The pH of water layer was adjusted to 4–6 by dropwise addition of NaOH (1 M), and the organic material was extracted with CH₂Cl₂ (50 mL). The aqueous layer was extracted with another portion of CH₂Cl₂ (50 mL). The combined organic layer was dried with anhydrous MgSO₄, evaporated under reduced pressure and purified by flash column chromatography using 50% EtOAc in hexanes to give **6e** as a colorless oil.

Acylation of 6a–e using symmetric anhydrides (7a–e). The symmetric anhydride of Boc-Phe was prepared in situ by mixing Boc-Phe (860 mg, 4 mmol) with EDCI (0.422 g, 2.2 mmol) in CH₂Cl₂ (10 mL, 0.2 M) for 5 min. To the solution of symmetric anhydride was added **6a–e** (1 mmol), and the reaction mixture was stirred for 24 h at room temperature. After dilution with CH₂Cl₂ (50 mL), the organic layer was washed with brine (50 mL), dried with anhydrous MgSO₄, and evaporated under reduced pressure. The residue was purified by flash column chromatography to obtain an Hcnb-linked dipeptide (**7a–e**).

Synthesis of Hcna

Methyl (E)-4-(2-(dimethylamino)vinyl)-3,5-dinitrobenzoate (13). To a stirred mixture of 4-Methyl-3,5-dinitrobenzoic acid (**12**) (2.0 g, 8.8 mmol) in anhydrous toluene (29 mL) was added DMF dimethyl acetal (3.2 mL, 24 mmol) dropwise at ambient temperature. The resulting mixture was heated to reflux in an oil bath and became dark red in color. After 16 h, the reaction was cooled to room temperature and the solvent removed under reduced pressure by co-evaporation with MeOH. The crude residue was recrystallized from hot ethanol to afford enamine **13** as a lustrous green solid which became red upon grinding.

Methyl 4-formyl-3,5-dinitrobenzoate (14). To a mixture of **13** (1.08 g, 3.66 mmol) and RuCl₃ solution (0.1 M in H₂O, 1.2 mL, 0.13 mmol) in CH₃CN–H₂O (6:1, 37 mL) was added NaIO₄ (2.0 g, 9.1 mmol) in portions at rt. The color quickly dissipated, and solids formed. The reaction continued until TLC analysis showed consumption of starting material (typically 10–30 min). After stirring for 30 min, the solids were removed by vacuum filtration and washed with CH₂Cl₂. The filtrate was partitioned between H₂O and CH₂Cl₂ and the aqueous layer was extracted twice with CH₂Cl₂. The combined organic extracts were washed with brine and sat. aq. Na₂S₂O₄, dried over Na₂SO₄, and concentrated under reduced pressure. The crude residue was

purified by flash column chromatography (30% EA/Hex) to afford **14** as an off-white solid.

Methyl 4-formyl-3-hydroxy-5-nitrobenzoate (15). A solution of **14** (678 mg, 2.67 mmol) in anhydrous DMF (5.1 mL) was added dropwise to a mixture of acetaldoxime [mixture of isomers] (325 μ L, 5.33 mmol) and K_2CO_3 (811 mg, 5.87 mmol) in anhydrous DMF (5.1 mL) dropwise at rt. The reaction mixture immediately turned bright red purple upon mixing, then brownish green after stirring overnight. After 21 h, H_2O (10 mL) was added to the reaction and the resulting solution extracted 5x with Et_2O . The aqueous layer was then acidified to pH ~1–2 with concentrated HCl and extracted 4x with CH_2Cl_2 . The combined organic extracts were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by flash chromatography (25% EA/Hex) to afford **15** as a bright yellow solid.

4-Formyl-3-hydroxy-5-nitrobenzoic acid (16). Solid $Ba(OH)_2 \cdot 8 H_2O$ (1.97 g, 6.24 mmol) was added in portions to a solution of **15** (468 mg, 2.08 mmol) in MeOH (23 mL) at rt. Following reaction completion (TLC), the mixture was concentrated under reduced pressure and the crude residue was partitioned between EtOAc and H_2O . The aqueous layer was extracted twice with EtOAc, acidified to pH ~1–2 with concentrated HCl and extracted with EtOAc (3x). The combined organic extracts were washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford **16** (379 mg, 86%) as a brown solid.

General procedure for solution-phase reductive amination (17a–c). HCl-AA-OtBu (0.666 mmol) was added to a solution of **15** (200.0 mg, 0.666 mmol) and DIEA (244 μ L, 1.40 mmol) in CH_3CN (9.5 mL). The reaction was stirred at rt for 45 min. Volatiles were removed and the residue was dissolved in MeOH (3.3 mL). $NaBH_4$ (75.6 mg, 2.00 mmol) was added, and the reaction was stirred for 15 min. Volatiles were removed and the residue was purified by flash column chromatography (20–45% EtOAc/Hex) to yield **17a–d**.

General procedure for solution-phase acylation (18a–c). A mixture of Fmoc-AA-OH (0.29 mmol) and DIPC (22.5 μ L, 0.14 mmol) was preactivated in dry CH_2Cl_2 :DMF (1:1) (0.6 mL) for 5 min. **17a–c** (0.13 mmol) as a solution in CH_2Cl_2 (2.1 mL) was added and the reaction was stirred for 2 h. The volatiles were removed and then dissolved in methanol (1.0 mL) and a solution of ammonium acetate (80.9 mg, 1.05 mmol) in water (0.3 mL) was added and the reaction was stirred overnight. Volatiles were removed and the residues were washed with water and extracted with ethyl acetate three times, concentrated, and then purified by flash column chromatography (30–40% EA/Hex).

Solid Phase Peptide Synthesis

Preparation of Hcna-loaded resin (22a–b). Fmoc-Sieber Amide PS Resin (0.6 mmol/g) or TentaGel S NH_2 (0.26 mmol/g) was swelled in DMF for 15 min, drained, and treated with piperidine-DMF (1:4, 1 $\times$ 2 mL, 30 min). The mixture was drained and washed with standard wash protocol [DMF (3 $\times$ 1 mL), CH_2Cl_2 (3 $\times$ 1 mL), MeOH (3 $\times$ 1 mL), CH_2Cl_2 (3 $\times$ 1 mL), then DMF (3 $\times$ 1 mL)] to afford a positive Kaiser ninhydrin test. To a mixture of *N*-Fmoc-6-aminoheptanoic acid (4 eq) and HATU (3.9 eq) in DMF (0.03 M) was added DIEA (8 eq) and this mixture was added to the resin (pre-washed with DMF) after a 5 min pre-activation period. After 2 h, the mixture was drained and washed with the standard wash protocol to afford a negative Kaiser ninhydrin test. The resin was then agitated with piperidine-DMF (1:4, 1 $\times$ 2 mL, 30 min). The mixture was drained, and the resin washed with standard wash protocol to afford a positive Kaiser ninhydrin test. To a mixture of Fmoc-Pro-OH (4 eq), HATU (3.9 eq) and HOAt (4 eq) in DMF (0.03 M) was added DIEA (8 eq) and this mixture was added to the resin (pre-washed with DMF) after a 5 min pre-activation period. After 2 h, the mixture was drained and washed with the standard wash protocol to afford a negative Kaiser ninhydrin test. The resin was then treated with piperidine-DMF (1:4, 1 $\times$ 2 mL, 30 min). The mixture was drained, and resin washed with standard wash protocol to afford a positive chloranil test. To a mixture of **16** (3 eq) and PyBOP in DMF (0.03 M) was added DIEA (6 eq) and this mixture was added to the resin (pre-washed with DMF) after a 5 min pre-activation period. After 2 h, the mixture was drained and washed with the standard wash protocol to afford a negative chloranil test.

General procedure for solid-phase reductive amination. The prepared resin **22a** or **22b** was swelled in DMF for 15 min then treated with a mixture of H-AA-ODMA or HCl-AA-OtBu (10 eq) and AcOH (10 eq) in CH_2Cl_2 (0.03 M). After 3 h, the mixture was drained and washed briefly with standard wash protocol then THF-MeOH (2:1). The resin was then taken up in THF-MeOH (2:1, 0.03 M) and treated with solid $NaBH_4$ (5 eq). After 2 h, the mixture was drained and washed with H_2O , standard wash protocol, then piperidine-DMF (1:4, 5 min). The resin was washed once more with standard wash protocol to afford a positive chloranil test.

General procedure for loading of second amino acid residue. A flame-dried flask was charged with the 2nd Amino Acid residue (4 eq) and dry CH_2Cl_2 (0.05 M), to which was added DIPC (2 eq) dropwise at rt. The mixture was stirred at rt for 12 min and the resulting solution was solubilized by addition of DMF (0.05 M). This solution was added directly to prepared resin **23a** or **23b** and the mixture

was agitated for 2 h, followed by the standard wash protocol to afford a slightly positive chloranil test. This procedure was repeated once more to afford resin-bound dipeptide (and a negative chloranil test).

General procedure for coupling of third amino acid residue. A solution of 1-octanthiol (10 eq), DIEA (5 eq), and DMF (0.03 M) was added to the prepared resin **24a** or **24b** and the mixture was agitated for 1 h, followed by the standard wash protocol. Next, a mixture of TIPS-OTf (10 eq), 2,6-Lutidine (10 eq), and DMF (0.03 M) was added to the resin and the mixture was agitated for 4 h, followed by the standard wash protocol. A flame-dried flask was charged with an Fmoc-amino acid (4 eq) and dry CH_2Cl_2 (0.05 M), to which was added DIPC (2 eq) dropwise at rt. The mixture was stirred at rt for 12 min and the resulting mixture was diluted with DMF (0.05 M).

The resin was agitated with a solution of DBU-1-Octanthiol-DMF (2:2:96, 1×2 mL, 3 min), washed with the standard wash protocol, and the pre-activated amino acid solution was added, taking care to minimize the time between deprotection and coupling. The mixture was agitated for 2 h, followed by standard wash protocol to afford a negative chloranil test. The resin was agitated with piperidine-DMF (1:4, 1×2 mL, 30 min). The mixture was drained, and the resin washed with standard wash protocol to afford a positive Kaiser ninhydrin test.

Fmoc half-cycle method. A solution of 1-octanthiol (10 eq), DIEA (5 eq), and DMF (0.03 M) was added to the prepared resin **24a** or **24b** and the mixture was agitated for 1 h, followed by the standard wash protocol. Next, a mixture of TIPS-OTf (10 eq), 2,6-Lutidine (10 eq), and DMF (0.03 M) was added to the resin and the mixture was agitated for 4 h, followed by the standard wash protocol. The C-terminal allyl protecting group was removed using $\text{Pd}(\text{PPh}_3)_4$ (0.25 eq), PhSiH_3 (25 eq) and DCM (3 mL) and agitating the reaction mixture for 2 h after which the following wash was done: DCM (X 3), DMF (X 3), 0.03 M sodium *N*, *N*-diethyldithiocarbamate in DMF (X 3), and DMF (X 3).

$\text{H-AA}_n\text{-OAllyl-CF}_3\text{COOH}$ (10 eq), HATU (10 eq), HOAt (10 eq) and DIEA (20 eq) in DMF (3 mL) was mixed and added to the reaction vessel and agitated for 4 h. The resin was washed using standard wash protocol.

General procedure for peptide cyclization (condition A). After complete synthesis of the desired Fmoc or Boc-protected peptide, the C-terminal DMA protecting group was removed using $\text{Pd}(\text{PPh}_3)_4$ (1 eq) and phenylsilane (20 eq) in CH_2Cl_2 under N_2 for 2 h, after which the resin was drained and washed with DMF (3×1 min), sodium *N*, *N*-diethyldithiocarbamate (0.03 M in DMF, 3×1 min), and CH_2Cl_2 (3×1 min). The resin was agitated with piperidine-DMF (1:4, 1×2 mL, 30 min). The mixture was drained, and

the resin washed with standard wash protocol to afford a positive Kaiser ninhydrin test. A mixture of PyAop (5 eq), HOAt (5 eq), and DIEA (10 eq) in DMF (0.03 M) was added directly to the resin. After 2 h, the mixture was drained and washed with the standard wash protocol to afford a negative Kaiser ninhydrin test.

General procedure for peptide cyclization (condition B). After complete synthesis of the desired Fmoc or Boc-protected peptide, the C-terminal DMA protecting group was removed using $\text{Pd}(\text{PPh}_3)_4$ (1 eq) and phenylsilane (20 eq) in CH_2Cl_2 under N_2 for 2 h, after which the resin was drained and washed with DMF (3×1 min), sodium *N*, *N*-diethyldithiocarbamate (0.03 M in DMF, 3×1 min), and CH_2Cl_2 (3×1 min). The resin was agitated with piperidine-DMF (1:4, 1×2 mL, 30 min). The mixture was drained, and the resin washed with standard wash protocol to afford a positive Kaiser ninhydrin test. A mixture of PyAOP (3 eq), 2,4,6-collidine (6 eq), and DIEA (6 eq) in 9:1 CH_2Cl_2 -DMF (0.03 M) was added directly to the resin. After 2 h, the mixture was drained and washed with the standard wash protocol to afford a negative Kaiser ninhydrin test.

General procedure for thioester synthesis. After synthesis of the desired Boc-protected peptide, the C-terminal DMA protecting group was removed using $\text{Pd}(\text{PPh}_3)_4$ (1 eq) and phenylsilane (20 eq) in CH_2Cl_2 under N_2 for 2 h, after which the resin was drained and washed with DMF (3×1 min), sodium *N*, *N*-diethyldithiocarbamate (0.03 M in DMF, 3×1 min), and CH_2Cl_2 (3×1 min). A solution of 1-octanthiol (10 eq), DIEA (10 eq), and DMF (0.03 M) was added to the resin and the mixture was agitated for 1 h, followed by the standard wash protocol. Next, a mixture of TIPS-OTf (10 eq), 2,6-Lutidine (10 eq), and DMF (0.03 M) were added to the resin and the mixture was agitated for 4 h, followed by the standard wash protocol. A solution of HATU (5 eq) and DIEA (20 eq) in DMF (0.03 M) was added to the resin and agitated for a 5 min period. Without draining, thiophenol (10 eq) was added to the mixture. After 2 h, the mixture was drained and washed with the standard wash protocol to afford a negative Kaiser ninhydrin test.

General Procedure for Sieber Amide cleavage. An aliquot (~1 mg) of resin-bound peptide was treated with 200 μL TFA/ H_2O / CH_2Cl_2 (2:1:97) and the mixture agitated for 5 min at rt. Following this, volatiles were removed by N_2 stream. The resulting residue was dissolved in 200 μL MeOH- H_2O (1:1), filtered and analyzed by UPLC-MS. Note: Product/DKP ratio was determined by integrating 214 and 254 nm peaks and are reported as relative percentages.

General Procedure for photocleavage. The resin-bound peptide was suspended in a 5 mL solution of 1:9 MeOH: CH_2Cl_2 in a fused quartz tube and agitated under 365 nm UV light for 1 h. The solvent was filtered to remove the resin and removed under reduced pressure. The resulting

residue was dissolved in MeOH-H₂O (1:1), filtered and analyzed by UPLC-MS and HPLC.

Results and Discussion

Synthesis of linker-anchored Dipeptides

Using the precursor molecules **1–5** (Fig. 2), each linker was covalently attached to glycine methyl ester to afford an anchored glycine residue (**6a–e**), which was then acylated with Boc-*L*-phenylalanine to afford an anchored dipeptide **7a–e** (Table 1). Each anchored dipeptide was photolyzed at 300 nm to afford Boc-Phe-Gly-OMe (**8**).

The linker precursors **1–5** were either commercially available or prepared using literature procedures (Stowell 1996; Vishwanathan 2006). Linkers **1** and **2** were anchored to methyl glycinate via nucleophilic displacement, whereas **3** and **4** employed the Amadori rearrangement (Amadori 1925; Isbell 1958) of an α -hydroxyketone, and **5** was attached to methyl glycinate through reductive amination. In all cases, acylation with the symmetric anhydride of Boc-*L*-Phe was found to afford the highest yield of anchored dipeptide. A survey of conditions for photolytic cleavage of the dipeptide found that photoexcitation at 300 nm in either methanol or chloroform, depending on the motif, afforded the fastest and cleanest reactions, but only the Dmp (**7d**) and Nve-linked dipeptides showed acceptable photolytic reactivity. Problems arose, however, when amino acids larger than glycine were anchored to Dmp and Nve. Both the initial attachment and subsequent acylation of those anchored amino acids proved to be quite problematic and afforded anchored dipeptides in modest yields at best.

Table 1 Photolabile linker candidate and their performance in amination, acylation and photolysis

$1-5 \xrightarrow{\text{GlyOMe HCl}} \begin{matrix} R \\ \\ \text{HN} \\ \\ \text{CO}_2\text{Me} \end{matrix} \xrightarrow[\text{CH}_2\text{Cl}_2]{(\text{Boc-Phe})_2\text{O}} \begin{matrix} \text{Ph} \\ \\ \text{BocHN} \\ \\ \text{C} \\ \\ \text{N} \\ \\ \text{CO}_2\text{Me} \end{matrix} \xrightarrow{300 \text{ nm}} 8$			
	6a , R = Op 6b , R = Mop 6c , R = Obz 6d , R = Dmp 6e , R = Nve	7a , R = Op 7b , R = Mop 7c , R = Obz 7d , R = Dmp 7e , R = Nve	
Starting material ^a	Amination (yield, %)	Acylation (yield, %)	Photolysis (yield, %)
1	6a (63) ^b	7a (81)	5 ^c
2	6b (78) ^{b, d}	7b (75)	23 ^c
3	6c (74) ^{c, e}	7c (50)	12 ^f
4	6d (85) ^g	7d (81)	75 ^f
5	6e (82) ^h	7e (77)	80 ^c

^aDetails of the preparation of **1–5** are given in the Supporting Information. ^bCH₃CN used as solvent. ^cMeOH used as solvent. ^dReaction run at 82 °C. ^eReaction run at 65 °C. ^fCHCl₃ used as solvent. ^gReaction run at 110 °C in PhCH₃. ^hNaBH₄ added to the reaction


In summary, Op, Mop, and Obz failed to release the dipeptide product efficiently, regardless of solvent and wavelength used, and so were discarded as linker candidates. Both the Nve and Dmp motifs displayed acceptable photolytic product release with the appropriate choices of solvent and irradiation wavelength; however, in both cases the modest yields for amino acid anchoring and acylation were insufficient for our needs, necessitating further modification to improve both steps of the reaction sequence.

Development of the second-generation Photolabile Linker

On the basis of both acylation and photocleavage yields the *o*-nitrobenzyl motif of the Nve linker was chosen as the basic skeleton for a second-generation linker, and the transacylation motif of the 6-hydroxy-4-methoxybenzyl (Hmb) (Sheppard 1993) and 6-hydroxynitrobenzyl (Hnb) (Smythe 1999) auxiliaries was incorporated into the linker design to improve acylation efficiency. The acid-labile Hmb auxiliary was devised by Sheppard for the protection of peptide amide backbones during the synthesis of ‘difficult’ peptide sequences. The related, photolabile Hnb auxiliary was developed by Smythe for the covalent modification of ‘difficult’ peptide sequences to facilitate their cyclization. Both the Hmb and Hnb auxiliaries employ an *ortho*-hydroxyl group to acylate bulky amino acids via esterification and subsequent O-to-N transacylation.

It was reasoned that the addition of an *o*-hydroxy substituent to the *o*-nitrobenzyl skeleton would assist both the acylation of the first amino acid and later photolytic cleavage. Attachment of a *para*-carboxylate to the *o*-nitrobenzyl skeleton provides a tether for covalent attachment of the linker to a spacer or resin and affords additional electron withdrawal to assist both the transacylation and photolysis reactions. Thus, the 2-hydroxy-4-carboxy-6-nitrobenzyl (Hcnb) linker emerged as a second-generation candidate for a photolabile BAL linker, as exemplified by the Hcnb-glycine adduct **9** (Fig. 3), derived from reductive amination of the aldehyde precursor 2-hydroxy-4-carboxy-6-nitrobenzaldehyde (Hcna) with glycine methyl ester. Treatment of **9** with a symmetric anhydride (Fig. 3) can result in the formation of the desired dipeptide **11** either through direct N-acylation or via initial acylation of the less hindered phenol followed by subsequent O-to-N acyl transfer. This alternate pathway was envisaged as a means to overcome the limitations of the Nve motif when attached to sterically hindered amino acids.

Fig. 3 Transacylation motif in the Hcnb linker



Synthesis of Hcna Linker

4-formyl-3-hydroxy-5-nitrobenzoic acid (**16**) was prepared from commercially available 4-Methyl-3,5-dinitrobenzoic acid (**12**) (Fig. 4). A condensation reaction between **12** and DMF-DMA afforded the previously reported (Starosotnikov 2005) enamine **13** in 72% yield. Ruthenium (III) chloride-catalyzed oxidative cleavage of the enamine using sodium metaperiodate generated the aldehyde **14** in 61% yield. Unilateral nucleophilic aromatic substitution of a single nitro-group using acetaldoxime to install the *ortho*-phenol group afforded ester-protected linker precursor **15** in 68% yield, which was used for solution-phase studies. Finally, hydrolysis of the methyl ester group using barium hydroxide afforded the free carboxylic acid **16** in 90% yield.

Solution-phase Synthesis of the Dipeptides Using Hcnb

We envisioned linking the peptide to the linker via a reductive amination between the primary amine of the first amino acid and the aldehyde precursor, in similar fashion to the loading protocol for the original BAL Linker. The resulting secondary amine can then be acylated with the next amino acid in sequence, forming a tertiary amide linkage between the peptide and linker. The presence of the *ortho*-phenol in

Hcnb is essential for complete acylation of the secondary amine, especially when utilizing amino acids with bulky side chains. Finally, photolysis in the short-wave UV range should cause a clean release of the dipeptide from the linker.

A series of dipeptides was synthesized in solution on the methyl ester-protected Hcna linker precursor **15** (Table 2). In solution, addition of the desired C-terminally protected amino acid to the aldehyde handle in the presence of DIEA in CH_3CN forms the imine intermediate. Reduction of this intermediate by sodium borohydride in methanol yields the Hcnb-linked amino acids **17a-c**. Acylation of the secondary amine with a pre-formed symmetric anhydride of the second amino acid resulted in acylation to give Hcnb-linked dipeptides, even in the case of bulky side-chains such as Thr. Photolysis at 365 nm in MeOH releases the desired dipeptides from the linker to give **19a-c**. Due to rotomers present in peptides attached to Hcnb, yields were reported over 2 steps.

Preparation of the resin-bound Hcna Linker

The free acid form of the Hcna linker, **16** (Fig. 4), was covalently anchored to aminomethyl-functionalized Tentagel resin **21a**, or Sieber TG resin **21b** via acylation of a resin-bound amine. A 6-amino-hexanoic acid (Ahx) spacer was employed with both solid supports to improve reactivity.




Fig. 4 Preparation of Hnca linker **16**

Table 2 Solution phase synthesis of dipeptides using Hnca



AA ₁	Amination, yield (%)	AA ₂	Acylation and Photocleavage, yield (%)
Val (17a)	51	Ala (18a & 19a)	38 (over 2 steps)
Phe (17b)	63	Val (18b & 19b)	35 (over 2 steps)
Lys(Z) (17c)	69	Leu (18c & 19c)	54 (over 2 steps)

Fig. 5 Preparation of resin-bound Hcna linkers **22a-b****Fig. 6** Addition of first and second amino acid residues to the Hcna linker

Additionally, a proline residue was introduced to reduce cross-linking between the amine and the aldehyde of the Hcna linker. In both cases, acylation of resin-bound amines was accomplished by coupling **16** to the free amine using either PyBOP or DCC (Fig. 5). Subsequent reactions of **22a** were monitored by photolytic cleavage from the linker and analysis of the crude product, whereas reactions of **22b** were monitored by cleavage using dilute acid and analysis of the crude product. Because of the time required for photolysis of Hcnb-linked peptides (1–2 h), reaction conditions were optimized using **22b** and then applied to **22a**.


Loading the Dipeptide by Reductive Amination Followed by Acylation

The first residue was anchored to the resin-bound Hcna linker via a 2-step reductive amination under acidic conditions. (Fig. 6). Quantitative acylation of the resin-bound amine was accomplished using a symmetric anhydride of the second residue (4 eq, 2 × 2 h) as determined by LC-MS analysis.

Loading of the Third Amino Acid


One of the major challenges with the Backbone Amide Linker concept is the formation of a diketopiperazine byproduct upon deprotection of the Fmoc group at the dipeptide stage. The most common method for overcoming this problem is to employ a sterically hindered C-terminal protecting group such as *t*-butyl or 1,1-dimethylallyl (DMA) (Hostetler 2018), thereby minimizing the formation of this undesired byproduct. However, it was found that quantitative DKP

formation was still obtained when using the Hcnb linker, regardless of the C-terminal protecting groups used. It was hypothesized that this results from Brønsted acid activation of the C-terminus by the nearby phenol proton, thereby catalyzing the formation of DKP. It was reasoned that this effect could be eliminated through removal of the phenol proton, thus avoiding Brønsted acid activation. The phenol group is required for efficient loading of the second amino acid residue, as well as for photocleavage of the final peptide, and so a method for temporary protection of the hydroxyl is required. Following the loading of the secondary amine, the phenol remains acylated with an excess of the second amino acid, until it is removed by the strongly nucleophilic piperidine upon deprotection of the N-terminal Fmoc group. We theorized that replacing piperidine with DBU, a relatively non-nucleophilic alternative for Fmoc deprotection, would leave the acylated phenol intact and minimize formation of the DKP byproduct. However, it was found that rapid, quantitative DKP formation was still obtained even when using DBU. Other methods were considered for blocking the phenol from interacting with the C-terminus, including silylation. To do this, the phenol was deacylated using 1-octanethiol, as indicated by our earlier experiences with Fmoc deprotection. However, due to the electronics of the ring, silylation of the phenol results in an exceedingly unstable silyl ether. Even attempts to observe the presence of a silyl group by LC-MS analysis proved unsuccessful. The most successful protection conditions used TIPS-OTf and 2,6-Lutidine for TIPS protection of the phenol on the linker. To monitor the effectiveness of silylation, the dipeptide N-terminus was deprotected for 3 min using a cocktail of 2% v/v DBU and 2% v/v 1-Octanethiol in DMF, followed

Table 3 DKP formation during coupling of the third residue (**27a-g**)

Dipeptide	Tripeptide	Tripeptide/DKP
Fmoc-Ala-*Val-ODMA	Fmoc-Gly-Ala-*Val-ODMA (27a)	100/0
Fmoc-Val-*Phe-ODMA	Fmoc-Phe-Val-*Phe-ODMA (27b)	96/6
Fmoc-Phe-*Lys(Boc)-ODMA	Fmoc-Val-Phe-*Lys(Boc)-ODMA (27c)	100/0
Fmoc-Ala-*Glu(tBu)-ODMA	Fmoc-Phe-Ala-*Glu(tBu)-ODMA (27d)	98/2
Fmoc-Val-*Leu-OrBu	Fmoc-Gly-Val-*Leu-OrBu (27e)	100/0
Fmoc-Ala-*Phe-OrBu	Fmoc-Phe-Ala-*Phe-OrBu (27f)	100/0
Fmoc-Ala-*Ala-OrBu	Fmoc-Leu-Ala-*Ala-OrBu (27g)	88/12

*residue anchored to Hcnb linker

**Fig. 7** The Fmoc-half cycle to synthesize tripeptides and avoid diketopiperazine formation

by immediate introduction of the third amino acid as a pre-formed symmetric anhydride. Various tripeptide sequences were tested to determine the efficacy of the TIPS group for reducing DKP formation (Table 3). In most cases, only trace amounts of DKP were observed by LC-MS, if any. The worst sequence tested was the Ala-Ala dipeptide, resulting in 12% DKP formation.

To completely avoid diketopiperazine formation, we followed the method developed by the Barany group, also known as the Fmoc-half cycle. In an Fmoc-half cycle, the first amino acid used in the reductive amination step is an allyl ester. This is followed by the acylation of the second amino acid. Instead of attaching the third amino acid to the N-terminus, the C-terminal allyl ester is deprotected and the

final amino acid is attached to the C-terminus by a peptide coupling reaction in which the C-terminal carboxylate is activated. In this case, the allyl ester was deprotected with catalytic Pd(PPh₃)₄ and excess PhSiH₃, and then coupled with another amino acid allyl ester through the C-terminus (Fig. 7). Then the rest of the peptide was built in N-to-C fashion by Fmoc SPPS until the peptide of desired length was built.

When using the Fmoc-half cycle method, we used allyl ester as the C-terminal protecting group. This was done to avoid an unwanted side reaction between we observed between C-terminal DMA esters and the residual 1-octanethiol remaining after Fmoc deprotection.




Fig. 8 Cyclization of 5-, 10-, and 14- residue cyclic peptides

Table 4 Structures and yields of peptide thioesters

Structure	Purity, %	Cleavage Yield, % ^a
H-Phe-Lys-Ala-Ala-Leu-SPh (31a)	99	83
H-Ala-Glu-Phe-Leu-Phe-SPh (31b)	99	66
H-Ala-Lys-Phe-Leu-Glu-SPh (31c)	75	71
H-Phe-Glu-Ala-Leu-Ala-SPh (31d)	99	92

^aYield was determined by HPLC analysis of the crude cleavage mixture

The Choice of Model Cyclic Peptides

The first peptide chosen was a short-chain analogue of the natural product somatostatin, cyclo[(D)-Trp-Lys-Gly-(β)-Ala-Phe] (**28**). The synthesis was carried out according to the procedures mentioned previously. The first residue H-Phe-ODMA was loaded onto the prepared resin **22a** via reductive amination. The subsequent linear synthesis was

then carried out as previously discussed. The head-to-tail cyclization was accomplished on the resin using PyAOP. Photocleavage from the resin at 365 nm for 1 h yielded the final cyclic pentapeptide **28** in 50% purity and 48% yield based on loading level. Two other previously synthesized cyclic peptides were chosen to examine the performance of the Hcnb linker on solid support. The cyclic decapeptide c-[Arg-(D)-Phe-Pro-Glu-Asp-Asn-Tyr-Glu-Ala-Ala, **29**] was synthesized to demonstrate the efficacy of the original Backbone Amide Linker, published by Albericio and Barany. The 14-residue cyclic peptide c-[Pro-Asp-Gly-Arg-Cys-Thr-Lys-Ser-Ile-Pro-Pro-Ile-Cys-Phe, **30**], also known as the Sunflower Trypsin Inhibitor, is a well-known natural product. The 14-mer cyclic Sunflower Trypsin Inhibitor poses some interesting synthetic challenges due to the high prevalence of β -branched amino acids and proline residues, which make selection of a cyclization point difficult. After several attempts, it was found that reductive amination of Phe onto the linker as the first residue, followed by cyclization at Pro yielded the best results. Cyclization and subsequent photocleavage from the resin yielded the final peptide in 90% purity and 80% yield based on loading level. The synthesis of the cyclic decapeptide was accomplished through reductive amination of Ala as the first residue and the subsequent linear synthesis. Following cyclization and photocleavage from the resin, the peptide was obtained in 95% purity and 90% yield based on loading level.

Synthesis of C-terminal Thioesters

The synthesis of C-terminal thioesters is an area of interest for their use in native chemical ligation (NCL) reactions. The Hcnb linker was used to demonstrate the on-resin synthesis of thioesters in high purities. It was established by solution-phase studies that the presence of a free phenol on the linker was problematic for synthesis of C-terminal thioesters, and so the phenol was TIPS protected prior to optimizing the coupling conditions for thioester formation. Use of carbodiimides as a coupling reagent yielded low conversions (0–30%). However, switching to HATU as a coupling reagent and preactivating the carboxylic acid for 5 min prior to addition of the thiol yielded complete conversion to the desired thioester in 2 h. A series of pentapeptides were synthesized to examine the scope of C-terminal residue compatibility with thioester synthesis. It was found that β -branched amino acids yield poor conversion to the thioester. However, it was reported by Kent that these sterically hindered amino acids are not amenable to NCL due to the steric hindrance of the carbonyl required for the transthioesterification and subsequent acyl transfer. The linear sequences were synthesized according to the standard procedure outlined previously. The final amino acid in the sequence was coupled as an N-Boc-protected residue. The C-terminal DMA protecting

group was removed using Pd(0) and phenylsilane. Following TIPS protection of the linker phenol, the C-terminus was pre-activated for 5 min with HATU and Hunig's base in DMF. Addition of thiophenol resulted in 95–100% conversion of carboxylic acid to thioester in 2 h. The thioester peptides were then globally deprotected using 5% H₂O in TFA for 1 h and cleaved from the resin using 365 nm UV irradiation for 1 h. Depending on the identity of the first amino acid residue, peptide thioesters were obtained in 75–99% purity and 66–92% yield as judged from HPLC analysis of the crude cleavage mixture (Table 3). No epimerization of the C-terminal residue was observed.

Conclusions

In a quest for a photolabile backbone amide linker for the solid-phase synthesis of cyclic and C-terminally modified peptides, a survey of various photolabile motifs was conducted, establishing the *o*-nitrobenzyl motif as the best candidate. Problems associated with many of the candidate motifs including the nitroveratryl linker led to the development of the 2-hydroxy-4-carboxy-6-nitrobenzyl (Hcnb) linker. A convenient and scalable synthesis of a benzaldehyde precursor (**16**) was developed, and general conditions were found for the anchoring of various amino esters to the linker and their acylation to form anchored dipeptides in good yield. Photolysis of the various Hcnb-linked dipeptides (**18a–c**) at 365 nm cleanly deprotected them in good yield in all cases. The linker can be attached to a wide variety of amino acids, can be selectively removed in the presence of most commonly used protecting groups and tolerates treatment with acid, base and Pd⁰. It was found that the Hcnb linker could be conveniently attached to an aminoethyl TG resin using a prolyl-6-aminohexanoic acid spacer, permitting its use in solid phase peptide synthesis. The ability of the Hcnb linker to facilitate on-resin, head-to-tail cyclization of peptides was demonstrated through the synthesis of three cyclic peptides. The utility of Hcnb to synthesize C-terminal thioesters was also demonstrated. We believe that this new photolabile linker should provide a valuable tool for the synthesis of cyclic and modified peptides, depsiptides and peptidomimetics such as oligoureas (Boeijen 1999; Boeijen 2001).

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10989-025-10752-9>.

Author Contributions K. K., M.N. and M.L. wrote the main manuscript text. SSK, M.H., M.N. and K.K. prepared the figures. All authors reviewed the manuscript.

Data Availability No datasets were generated or analysed during the

current study.

Declarations

Competing Interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Amadori M (1925) Atti accad. Naz Lincei 2(6):337
- Bellof D, Mutter M (1985) A new phenacyl-type handle for polymer supported peptide synthesis. *Chimia* 39:317
- Bergmark WR, Barnes C, Clark J, Paparian S, Marynowski S (1985) Photoenolization with α -chloro substituents. *J Org Chem* 50(26):5612–5615. <https://doi.org/10.1021/jo00350a036>
- Biron E, Chatterjee J, Ovadia O, Langenegger D, Bruegggen J, Hoyer D, Schmid HA, Jelinek R, Gilon C, Hoffman A, Kessler H (2008) Improving oral bioavailability of peptides by multiple N-methylation: somatostatin analogues. *Angew Chem Int Ed* 47:2595. <http://doi.org/10.1002/anie.200705797>
- Boas U, Brask J, Christensen JB, Jensen KJ (2002) The ortho backbone amide linker (o-BAL) is an easily prepared and highly acid-labile handle for solid-phase synthesis. *J Comb Chem* 4:223. <http://doi.org/10.1021/cc010070h>
- Boas U, Christensen JB, Jensen KJ (2004) Two dialkoxynaphthalene aldehydes as backbone amide linkers for solid-phase synthesis. *J Comb Chem* 6:497. <https://doi.org/10.1021/cc034056b>
- Boeijen A, Liskamp RM (1999) Solid-Phase Synthesis of Oligoureia Peptidomimetics. *J. Eur. J. Org. Chem.* [https://doi.org/10.1002/\(SICI\)1099-0690\(199909\)1999:9<2127::AID-EJOC2127>3.0.CO;2-T](https://doi.org/10.1002/(SICI)1099-0690(199909)1999:9<2127::AID-EJOC2127>3.0.CO;2-T)
- Boeijen A, van Ameijde J, Liskamp RMJ (2001) Solid-Phase synthesis of oligoureia peptidomimetics employing the Fmoc protection strategy. *J Org Chem* 66:8454. <https://doi.org/10.1021/jo010656q>
- Bourne GT, Golding SW, Meutermans WDF, Smythe ML (2001) Synthesis of a cyclic peptide library based on the somatostatin sequence using the backbone amide linker approach. *Lett Pept Sci* 7:311. <https://doi.org/10.1023/A:1013057832044>
- Bourne GT, Meutermans WDF, Smythe ML (1999) The development of solid phase protocols for a backbone amide linker and its application to the Boc-based assembly of linear peptides tetrahedron Lett. 40:7271. [https://doi.org/10.1016/S0040-4039\(99\)01473-2](https://doi.org/10.1016/S0040-4039(99)01473-2)
- Brask J, Albericio F, Jensen KJ (2003) Fmoc solid-phase synthesis of peptide thioesters by masking as trithioortho esters. *Org Lett* 5:2951. <https://doi.org/10.1021/ol0351044>
- Coulouarn Y, Lihmann I, Jegou S, Anouar Y, Tostivint H, Beauvillain JC, Conlon JM, Bern HA, Vaudry A (1998) Cloning of the cDNA encoding the Urotensin II precursor in frog and human reveals intense expression of the Urotensin II gene in motoneurons of the spinal cord. *Proc Natl Acad Sci* 95:15803. <https://doi.org/10.1073/pnas.95.26.15803>
- Darkin-Rattray SJ, Gurnett AM, Myers RW, Dulski PM, Crumley TM, Allocco JJ, Cannova C, Meinke PT, Colletti SL, Bednarek MA, Singh SB, Goetz MA, Dombrowski AW, Polishook JD, Schmatz DM (1996) Apicidin: A novel antiprotozoal agent that inhibits parasite histone deacetylase. *Proc. Natl. Acad. Sci.* 93, 13143. <https://doi.org/10.1073/pnas.93.23.13143>
- Dawson PE, Muir TW, Clark-Lewis I, Kent SB (1994) Synthesis of proteins by native chemical ligation. *Science* 266:776. <https://doi.org/10.1126/science.7973629>
- Dougherty PG, Sahni A, Pei D (2019) Understanding cell penetration of cyclic peptides. *Chem Rev* 119:10241. <https://doi.org/10.1021/acs.chemrev.9b00008>
- Foister S, Taylor LL, Feng J, Chen W, Lin A, Cheng F, Smith AB III, Hirschmann R (2006) Design and synthesis of potent cystine-free Cyclic hexapeptide agonists at the human Urotensin receptor. *Org Lett* 8:1799. <https://doi.org/10.1021/ol060278h>
- Fusetani N, Matsunaga S (1993) Bioactive sponge peptides. *Chem Rev* 93:1793. <https://doi.org/10.1021/cr00021a007>
- Gallop MA, Barrett RW, Dower WJ, Fodor SPA, Gordon EM (1994) Applications of combinatorial technologies to drug discovery. 1. Background and peptide combinatorial libraries. *J Med Chem* 37:1233. <https://doi.org/10.1021/jm00035a001>
- Gause GF, Brazhnikova MG (1944) Gramicidin S and its use in the treatment of infected wounds. *Nature* 154:703. <https://doi.org/10.1038/154703a0>
- Guillaumie F, Kappel JC, Kelly NM, Barany G, Jensen KJ (2000) Solid-phase synthesis of C-terminal peptide aldehydes from amino acetals anchored to a backbone amide linker (BAL) handle. *Tetrahedron Lett* 41:6131. [https://doi.org/10.1016/S0040-4039\(00\)00950-3](https://doi.org/10.1016/S0040-4039(00)00950-3)
- Guillier F, Orain D, Bradley M (2000) Linkers and cleavage strategies in solid-phase organic synthesis and combinatorial chemistry. *Chem. Rev.* 100, 2091. <https://doi.org/10.1021/cr980040+>
- Heidler P, Link A (2005) N-acyl-N-alkyl-sulfonamide anchors derived from kenner's safety-catch linker: powerful tools in bioorganic and medicinal chemistry. *Bioorg Med Chem* 13:585. <https://doi.org/10.1016/j.bmc.2004.10.045>
- Hostettler MA, Lipton MA (2018) An optimized preparation of 1,1-dimethylallyl esters and their application to solid-phase peptide synthesis. *J Org Chem* 83(15):7762–7770. <https://doi.org/10.1021/acs.joc.8b00658>
- Humphrey JM, Chamberlin AR (1997) Chemical synthesis of natural product peptides: coupling methods for the incorporation of non-coded amino acids into peptides. *Chem Rev* 97:2243. <https://doi.org/10.1021/cr950005s>
- Isbell HS, Frush HL (1958) Mutarotation, hydrolysis, and rearrangement reactions of glycosylamines. *J Org Chem* 23:1309
- Jensen KJ, Alsina J, Songster MF, Vagner J, Albericio F, Barany G (1998) Backbone amide linker strategies for the solid-phase synthesis of C-terminal modified peptides. *J Am Chem Soc* 120:5441. <https://doi.org/10.1385/1-59259-877-3:195>
- Jessing M, Brandt M, Jensen KJ, Christensen JB, Boas U (2006) Thio-phenyl backbone amide linkers, a new class of easily prepared and highly acid-labile linkers for solid-phase synthesis. *J Org Chem* 71:6734. <https://doi.org/10.1021/jo060687r>
- Kang SS, Lipton MA (2008) Chap. 17 Photolabile and Miscellaneous Linkers/Resins in 'The Power of Functional Resins in Organic Synthesis'; Editor: Albericio, F.; Wiley-VCH; ISBN: 978-3527319367
- Kappel JC, Barany G (2005) Backbone amide linker (BAL) strategy for N⁹-9-fluorenylmethoxycarbonyl (Fmoc) solid-phase synthesis of peptide aldehydes. *J Pept Sci* 11:525. <https://doi.org/10.1002/psc.614>
- Katsara M, Tselios T, Deraos S, Deraos G, Matsoukas M, Lazoura E, Matsoukas J, Apostolopoulos V (2006) Design and synthesis of a cyclic double mutant peptide (cyclo(87–99)[A⁹¹,A⁹⁶]MBP_{87–99})

- induces altered responses in mice after conjugation to mannan: implications in the immunotherapy of multiple sclerosis. *Curr Med Chem* 13:2221. <https://doi.org/10.1021/jm801250v>
- Kenner GW, McDermott JR, Sheppard RC (1971) The safety catch principle in solid phase peptide synthesis. *J Chem Soc Chem Commun.* <https://doi.org/10.1039/C29710000636>
- Kessler H (1982) Conformation and biological activity of cyclic peptides. *Angew Chem Int Ed* 21:512. <https://doi.org/10.1002/anie.198205121>
- Kessler M, Glatthar R, Giese B, Bochet C (2003) Sequentially photocleavable protecting groups in solid-phase synthesis. *Org Lett* 5(8):1179–1181. <https://doi.org/10.1021/ol027454g>
- Krishnamoorthy R, Vazquez-Serrano LD, Turk JA, Kowalski JA, Benson AG, Breau NT, Lipton MA (2006) Complete stereochemistry of neamphamide A and absolute configuration of the β -Methoxytyrosine residue in Papuamide B. *J Am Chem Soc* 128:15392. <https://doi.org/10.1021/jo0508853>
- Liang X, Vezina-Dawod S, Bedard F, Porte K, Biron E (2016) One-pot photochemical ring-opening/cleavage approach for the synthesis and decoding of cyclic peptide libraries. *Org Lett* 18(5):1174–1177. <https://doi.org/10.1021/acs.orglett.6b00296>
- Li H, Dong S (2017) Recent advances in the preparation of Fmoc-SPPS-based peptide thioester and its surrogates for NCL-type reactions. *Sci China Chem* 60:201. <https://doi.org/10.1007/s11426-016-0381-1>
- Liley MJ, Johnson T, Gilson SE (2006) An improved aldehyde linker for the solid phase synthesis of hindered amides. *J Org Chem* 71:1322. <https://doi.org/10.1021/jo051742j>
- Lloyd-Williams P, Albericio F, Giralt E (1993) Solid-phase synthesis of peptides using allylic anchoring groups 2. Palladium-catalysed cleavage of Fmoc-protected peptides. *Tetrahedron* 49:11065. [https://doi.org/10.1016/S0040-4039\(00\)73378-8](https://doi.org/10.1016/S0040-4039(00)73378-8)
- McAllister LA, McCormick RA, Procter DJ (2005) Sulfide- and selenide-based linkers in phase tag-assisted synthesis. *Tetrahedron* 61:11527. <https://doi.org/10.1016/j.tet.2005.08.056>
- Mende F, Seitz O (2011) 9-Fluorenylmethoxycarbonyl-based solid-phase synthesis of peptide α -thioesters. *Angew Chem Int Ed* 50:1232. <https://doi.org/10.1002/anie.201005180>
- Mihara H, Yamabe S, Nildome T, Aoyagi H, Kumagai H (1995) Efficient preparation of cyclic peptide mixtures by solid phase synthesis and cyclization cleavage with oxime resin. *Tetrahedron Lett* 36:4837. [https://doi.org/10.1016/0040-4039\(95\)00906-S](https://doi.org/10.1016/0040-4039(95)00906-S)
- Mori A, Akahoshi I, Hashimoto M, Doi T, Takahashi T (2004) Combinatorial library synthesis of two- and three-ring benzenoid amides on solid support. *Tetrahedron Lett* 45:813. <https://doi.org/10.1016/j.tetlet.2003.11.043>
- Nilsson BL, Kiessling LL, Raines RT (2000) Staudinger ligation: a peptide from a thioester and azide. *Org. Lett.* 2, 1939. <https://doi.org/10.1021/ol0060174>
- Patchornik A, Amit B, Woodward RB (1970) Photosensitive protecting groups. *J Am Chem Soc* 92:6333. <https://doi.org/10.1021/ja00724a041>
- Peach JM, Pratt AJ, Snaith JS (1995) Photolabile benzoin and furoin esters of a biologically active peptide. *Tetrahedron* 51:10013. [https://doi.org/10.1016/0040-4020\(95\)00573-Q](https://doi.org/10.1016/0040-4020(95)00573-Q)
- Pittelkow M, Boas U, Christensen JB (2006) Carbocations in action. Design, synthesis, and evaluation of a highly acid-sensitive naphthalene-based backbone amide linker for solid-phase synthesis. *Org Lett* 8:5817. <https://doi.org/10.1021/ol062410j>
- Pittelkow M, Boas U, Jessing M, Jensen KJ, Christensen JB (2005) Role of the peri-effect in synthesis and reactivity of highly substituted naphthaldehydes: a novel backbone amide linker for solid-phase synthesis. *Org Biomol Chem* 3:508. <https://doi.org/10.1039/B412971G>
- Richter LS, Tom JYK, Brunier JP (1994) Peptide-cyclizations on solid support: a fast and efficient route to small cyclopeptides. *Tetrahedron Lett* 35:5547. [https://doi.org/10.1016/S0040-4039\(0\)77243-1](https://doi.org/10.1016/S0040-4039(0)77243-1)
- Rosenbaum C, Waldmann H (2001) Solid phase synthesis of cyclic peptides by oxidative cyclative cleavage of an aryl hydrazide linker—synthesis of stylostatin 1. *Tetrahedron Lett* 42:5677. [https://doi.org/10.1016/S0040-4039\(01\)01075-9](https://doi.org/10.1016/S0040-4039(01)01075-9)
- Samanen J, Ali F, Romoff T, Calvo R, Sorenson E, Vasko J, Storer B, Berry D, Bennett D, Strohsacker M, Powers D, Stadel J, Nichols A (1991) Development of a small RGD peptide fibrinogen receptor antagonist with potent antiaggregatory activity *in vitro*. *J Med Chem* 34:3114. <https://doi.org/10.1021/jm00114a022>
- Scott P, Steel P (2006) Diversity linker units for solid-phase organic synthesis. *Eur J Org Chem.* <https://doi.org/10.1002/ejoc.200500804>
- Sheehan JC, Davies GD (1964) Facile Alkyl-Oxygen Ester Cleavage. *J. Org. Chem.* 29, 2006
- Sheehan JC, Umezawa K (1973) Phenacyl photosensitive blocking groups. *J Org Chem* 38:3771
- Sheppard RC, Johnson T, Quibell M, Owen D (1993) A reversible protecting group for the amide bond in peptides. Use in the synthesis of 'difficult sequences'. *J Chem Soc Chem Commun.* <https://doi.org/10.1039/C39930000369>
- Shevach EM (1985) The effects of cyclosporin A on the immune system. *Ann Rev Immuno* 3:397. <https://doi.org/10.1146/annurev.iy.03.040185.002145>
- Smythe M, Meutermaans W, Golding S, Bourne G, Miranda L, Dooley M, Alewood P (1999) Synthesis of difficult cyclic peptides by inclusion of a novel photolabile auxiliary in a ring contraction strategy. *J Am Chem Soc* 121:9790. <https://doi.org/10.1021/ja992173y>
- Soural M, Krchnak V (2007) Efficient solid-phase synthesis of 2-substituted-3-hydroxy-4(1H)-quinolinone-7-carboxamides with two diversity positions. *J Comb Chem* 9:793. <https://doi.org/10.1021/cc070071v>
- Springer J, de Cuba KR, Calvet-Vitale S, Geenevasen JAJ, Hermkens PHH, Hiemstra H, van Maarseveen JH (2008) Backbone amide linker strategy for the synthesis of 1,4-triazole-containing cyclic tetra- and pentapeptides. *Eur J Org Chem* 15:2592. <https://doi.org/10.1002/ejoc.200800143>
- Starosotnikov AM, Lobach AV, Shevelev SA (2005) An efficient One-Step method for the conversion of β (Dimethylamino) styrenes into arylacetonitriles. *Synthesis* 17:2830–2832. <https://doi.org/10.1055/s-2005-918405>
- Stowell MHB, Rock RS, Rees DC, Chan SI (1996) Efficient synthesis of photolabile alkoxy benzoin protecting groups. *Tetrahedron Lett* 37(3):307–310. [https://doi.org/10.1016/0040-4039\(95\)02159-0](https://doi.org/10.1016/0040-4039(95)02159-0)
- Veber DF, Freidinger RM, Perlow DS, Paleveda WJ, Holly FW, Strachan RG, Nutt RF, Arison BH, Homnick C, Randall WC, Glitzer MS, Saperstein R, Hirschmann R (1981) A super active cyclic hexapeptide analog of somatostatin. *Nature.* [https://doi.org/10.1016/0024-3205\(84\)90009-2](https://doi.org/10.1016/0024-3205(84)90009-2)
- Viswanathan CL, Chaudhari AS (2006) Synthesis and evaluation of uterine relaxant activity for a novel series of substituted p-hydroxyphenylethanolamines. *Bioorg Med Chem* 14(19):6581–6585. <https://doi.org/10.1016/j.bmc.2006.06.006>
- White TR, Renzelman CM, Rand AC, Rezaei T, McEwen CM, Gelev VM, Turner RA, Linington RG, Leung SS, Kalgutkar AS, Bauman JN, Zhang Y, Liras S, Price DA, Mathiowetz AM, Jacobson MP, Lokey RS (2011) On-resin N-methylation of cyclic peptides for discovery of orally bioavailable scaffolds. *Nat Chem Biol* 7:810. <https://doi.org/10.1038/nchembio.664>